

AR and DUSP9 are promising druggable targets for treating Hypertension that control activity of BCL6, ZBTB7B and ZNF462 transcription factor on of highly methylated genes

Demo User

geneXplain GmbH

info@genexplain.com

Data received on 08/04/2022 ; Run on 10/12/2023 ; Report generated on 10/12/2023

Genome Enhancer release 3.3 (TRANSFAC®, TRANSPATH® and HumanPSD™ release 2023.2)



Abstract

In the present study we applied the software package "Genome Enhancer" to a data set that contains *epigenomics* data. The study is done in the context of *Hypertension*. The goal of this pipeline is to identify potential drug targets in the molecular network that governs the studied pathological process. In the first step of analysis pipeline discovers transcription factors (TFs) that regulate genes activities in the pathological state. The activities of these TFs are controlled by so-called master regulators, which are identified in the second step of analysis. After a subsequent druggability checkup, the most promising master regulators are chosen as potential drug targets for the analyzed pathology. At the end the pipeline comes up with (a) a list of known drugs and (b) investigational active chemical compounds with the potential to interact with selected drug targets.

From the data set analyzed in this study, we found the following TFs to be potentially involved in the regulation of the highly methylated genes: BCL6, ZBTB7B and ZNF462. The subsequent network analysis suggested

- RSK
- FAM123B
- trkC
- MKP-4
- AR-isoform1

as the most promising molecular targets for further research, drug development and drug repurposing initiatives on the basis of identified molecular mechanism of the studied pathology. Having checked the actual druggability potential of the full list of identified targets, both, via information available in medical literature and via cheminformatics analysis of drug compounds, we have identified the following drugs as the most promising treatment candidates for the studied pathology: Imatinib, Flavopiridol, Moexipril and 2,5,7-Trihydroxynaphthoquinone.

1. Introduction

Recording "-omics" data to measure gene activities, protein expression or metabolic events is becoming a standard approach to characterize the pathological state of an affected organism or tissue. Increasingly, several of these methods are applied in a combined approach leading to large "multiomics" datasets. Still the challenge remains how to reveal the underlying molecular mechanisms that render a given pathological state different from the norm. The disease-causing mechanism can be described by a re-wiring of the cellular regulatory network, for instance as a result of a genetic or epigenetic alterations influencing the activity of relevant genes. Reconstruction of the disease-specific regulatory networks can help identify potential master regulators of the respective pathological process. Knowledge about these master regulators can point to ways how to block a pathological regulatory cascade. Suppression of certain molecular targets as components of these cascades may stop the pathological process and cure the disease.

Conventional approaches of statistical "-omics" data analysis provide only very limited information about the causes of the observed phenomena and therefore contribute little to the understanding of the pathological molecular mechanism. In contrast, the "upstream analysis" method [1-4] applied here has been devised to provide a casual interpretation of the data obtained for a pathology state. This approach comprises two major steps: (1) analysing promoters and enhancers of highly methylated genes for the transcription factors (TFs) involved in their regulation and, thus, important for the process under study; (2) re-constructing the signaling pathways that activate these TFs and identifying master regulators at the top of such pathways. For the first step, the database TRANSFAC® [6] is employed together with the TF binding site identification algorithms Match [7] and CMA [8]. The second step involves the signal transduction database TRANSPATH® [9] and special graph search algorithms [10] implemented in the software "Genome Enhancer".

The "upstream analysis" approach has now been extended by a third step that reveals known drugs suitable to inhibit (or activate) the identified molecular targets in the context of the disease under study. This step is performed by using information from HumanPSD™ database [5]. In addition, some known drugs and investigational active chemical compounds are subsequently predicted as potential ligands for the revealed molecular targets. They are predicted using a pre-computed database of spectra of biological activities of chemical compounds of a library of 2245 known drugs and investigational chemical compounds from HumanPSD™ database. The spectra of biological activities for these compounds are computed using the program PASS on the basis of a (Q)SAR approach [11-13]. These predictions can be used for the research purposes - for further drug development and drug repurposing initiatives.

2. Data

For this study the following experimental data was used:

Table 1. Experimental datasets used in the study

File name	Data type
df_hp	Epigenomics
df_norm	Epigenomics

Hypertension

 LCL_0001
 LCL_0005
 LCL_0016
 LCL_0019
 LCL_0021
 LCL_0023
 LCL_0028
 LCL_0030
 LCL_0032
 LCL_0037
 LCL_0040
 LCL_0041
 LCL_0044
 LCL_0048
 LCL_0050
 LCL_0052
 LCL_0054
 LCL_0055
 LCL_0057
 LCL_0060
 LCL_0063
 LCL_0072
 LCL_0082
 LCL_0084
 LCL_0085
 LCL_0089
 LCL_0092
 LCL_0093
 LCL_0101
 LCL_0105
 LCL_0111
 LCL_0112
 LCL_0113
 LCL_0120
 LCL_0125
 LCL_0131
 LCL_0141

Control

 LCL_0007
 LCL_0009
 LCL_0013
 LCL_0017
 LCL_0018
 LCL_0022
 LCL_0029
 LCL_0043
 LCL_0046
 LCL_0066
 LCL_0069
 LCL_0074
 LCL_0076
 LCL_0077
 LCL_0083
 LCL_0087
 LCL_0088
 LCL_0091
 LCL_0095
 LCL_0003
 LCL_0097
 LCL_0102
 LCL_0107
 LCL_0108
 LCL_0122
 LCL_0128
 LCL_0129
 LCL_0133
 LCL_0136
 LCL_0144
 LCL_0145
 LCL_0147
 LCL_0148
 LCL_0150
 LCL_0155
 LCL_0158
 LCL_0159

df_hp
E LCL_0162
df_hp
E LCL_0171
df_hp
E LCL_0175
df_hp
E LCL_0177
df_hp
E LCL_0189
df_hp
E LCL_0190
df_hp
E LCL_0192
df_hp
E LCL_0193
df_hp
E LCL_0194
df_hp
E LCL_0198
df_hp
E LCL_0199
df_hp
E LCL_0200
df_hp
E LCL_0205
df_hp
E LCL_0211
df_hp
E LCL_0213
df_hp
E LCL_0214
df_hp
E LCL_0217
df_hp
E LCL_0233
df_hp
E LCL_0235
df_hp
E LCL_0236
df_hp
E LCL_0237
df_hp
E LCL_0241
df_hp
E LCL_0246
df_hp
E LCL_0255
df_hp
E LCL_0256
df_hp
E LCL_0257
df_hp
E LCL_0262
df_hp
E LCL_0265
df_hp
E LCL_0266
df_hp
E LCL_0272
df_hp
E LCL_0280
df_hp
E LCL_0285
df_hp
E LCL_0298
df_hp
E LCL_0299
df_hp
E LCL_0310
df_hp
E LCL_0314
df_hp
E LCL_0322
df_hp
E LCL_0324
df_hp

df_norm
E LCL_0135
df_norm
E LCL_0164
df_norm
E LCL_0165
df_norm
E LCL_0167
df_norm
E LCL_0168
df_norm
E LCL_0174
df_norm
E LCL_0176
df_norm
E LCL_0179
df_norm
E LCL_0183
df_norm
E LCL_0188
df_norm
E LCL_0196
df_norm
E LCL_0197
df_norm
E LCL_0204
df_norm
E LCL_0209
df_norm
E LCL_0219
df_norm
E LCL_0220
df_norm
E LCL_0221
df_norm
E LCL_0223
df_norm
E LCL_0224
df_norm
E LCL_0232
df_norm
E LCL_0234
df_norm
E LCL_0244
df_norm
E LCL_0245
df_norm
E LCL_0253
df_norm
E LCL_0258
df_norm
E LCL_0260
df_norm
E LCL_0264
df_norm
E LCL_0274
df_norm
E LCL_0281
df_norm
E LCL_0290
df_norm
E LCL_0291
df_norm
E LCL_0293
df_norm
E LCL_0294
df_norm
E LCL_0295
df_norm
E LCL_0301
df_norm
E LCL_0303
df_norm
E LCL_0306
df_norm
E LCL_0307
df_norm

LCL_0327
df_hp
LCL_0338
df_hp
LCL_0350
df_hp
LCL_0351
df_hp
LCL_0352
df_hp
LCL_0355
df_hp
LCL_0356
df_hp
LCL_0363
df_hp
LCL_0366
df_hp
LCL_0369
df_hp
LCL_0372
df_hp
LCL_0374
df_hp
LCL_0392
df_hp
LCL_0393
df_hp
LCL_0398
df_hp
LCL_0399
df_hp
LCL_0401
df_hp
LCL_0403
df_hp
LCL_0406
df_hp
LCL_0408
df_hp
LCL_0410
df_hp
LCL_0417
df_hp
LCL_0420
df_hp
LCL_0430
df_hp
LCL_0434
df_hp
LCL_0441
df_hp
LCL_0444
df_hp
LCL_0445
df_hp
LCL_0447
df_hp
LCL_0460
df_hp
LCL_0468
df_hp
LCL_0473
df_hp
LCL_0478
df_hp
LCL_0483
df_hp
LCL_0484
df_hp
LCL_0486
df_hp
LCL_0507
df_hp
LCL_0509
df_hp

LCL_0315
df_norm
LCL_0317
df_norm
LCL_0318
df_norm
LCL_0319
df_norm
LCL_0325
df_norm
LCL_0321
df_norm
LCL_0329
df_norm
LCL_0336
df_norm
LCL_0337
df_norm
LCL_0341
df_norm
LCL_0348
df_norm
LCL_0361
df_norm
LCL_0365
df_norm
LCL_0375
df_norm
LCL_0388
df_norm
LCL_0419
df_norm
LCL_0427
df_norm
LCL_0428
df_norm
LCL_0429
df_norm
LCL_0437
df_norm
LCL_0442
df_norm
LCL_0456
df_norm
LCL_0476
df_norm
LCL_0496
df_norm
LCL_0498
df_norm
LCL_0499
df_norm
LCL_0502
df_norm
LCL_0503
df_norm
LCL_0504
df_norm
LCL_0505
df_norm
LCL_0506
df_norm
LCL_0531
df_norm
LCL_0533
df_norm
LCL_0538
df_norm
LCL_0568
df_norm
LCL_0578
df_norm
LCL_0585
df_norm
LCL_0586
df_norm

LCL_0511
df_hp
LCL_0512
df_hp
LCL_0514
df_hp
LCL_0515
df_hp
LCL_0518
df_hp
LCL_0519
df_hp
LCL_0522
df_hp
LCL_0528
df_hp
LCL_0532
df_hp
LCL_0534
df_hp
LCL_0537
df_hp
LCL_0547
df_hp
LCL_0552
df_hp
LCL_0554
df_hp
LCL_0542
df_hp
LCL_0569
df_hp
LCL_0570
df_hp
LCL_0571
df_hp
LCL_0581
df_hp
LCL_0584
df_hp
LCL_0590
df_hp
LCL_0597
df_hp
LCL_0604
df_hp
LCL_0610
df_hp
LCL_0611
df_hp
LCL_0612
df_hp
LCL_0620
df_hp
LCL_0630
df_hp
LCL_0631
df_hp
LCL_0637
df_hp
LCL_0641
df_hp
LCL_0642
df_hp
LCL_0659
df_hp
LCL_0660
df_hp
LCL_0665
df_hp
LCL_0673
df_hp
LCL_0677
df_hp
LCL_0678
df_hp
LCL_0687

LCL_0596
df_norm
LCL_0599
df_norm
LCL_0605
df_norm
LCL_0607
df_norm
LCL_0616
df_norm
LCL_0618
df_norm
LCL_0625
df_norm
LCL_0626
df_norm
LCL_0627
df_norm
LCL_0633
df_norm
LCL_0634
df_norm
LCL_0638
df_norm
LCL_0640
df_norm
LCL_0643
df_norm
LCL_0644
df_norm
LCL_0645
df_norm
LCL_0655
df_norm
LCL_0663
df_norm
LCL_0672
df_norm
LCL_0676
df_norm
LCL_0690
df_norm
LCL_0692
df_norm
LCL_0693
df_norm
LCL_0700
df_norm
LCL_0705
df_norm
LCL_0709
df_norm
LCL_0717
df_norm
LCL_0719
df_norm
LCL_0718
df_norm
LCL_0724
df_norm
LCL_0728
df_norm
LCL_0731
df_norm
LCL_0734
df_norm
LCL_0738
df_norm
LCL_0739
df_norm
LCL_0745
df_norm
LCL_0749
df_norm
LCL_0754
df_norm
LCL_0756

df_hp
LCL_0691
df_hp
LCL_0694
df_hp
LCL_0697
df_hp
LCL_0701
df_hp
LCL_0702
df_hp
LCL_0707
df_hp
LCL_0711
df_hp
LCL_0723
df_hp
LCL_0726
df_hp
LCL_0727
df_hp
LCL_0732
df_hp
LCL_0733
df_hp
LCL_0737
df_hp
LCL_0741
df_hp
LCL_0742
df_hp
LCL_0743
df_hp
LCL_0755
df_hp
LCL_0773
df_hp
LCL_0774
df_hp
LCL_0775
df_hp
LCL_0778
df_hp
LCL_0783
df_hp
LCL_0785
df_hp
LCL_0795
df_hp
LCL_0796
df_hp
LCL_0800
df_hp
LCL_0808
df_hp
LCL_0810
df_hp
LCL_0814
df_hp
LCL_0817
df_hp
LCL_0836
df_hp
LCL_0837
df_hp
LCL_0866
df_hp
LCL_0869
df_hp
LCL_0871
df_hp
LCL_0873
df_hp
LCL_0877
df_hp
LCL_0893
df_hp

df_norm
LCL_0757
df_norm
LCL_0762
df_norm
LCL_0764
df_norm
LCL_0765
df_norm
LCL_0769
df_norm
LCL_0780
df_norm
LCL_0784
df_norm
LCL_0788
df_norm
LCL_0789
df_norm
LCL_0793
df_norm
LCL_0797
df_norm
LCL_0799
df_norm
LCL_0805
df_norm
LCL_0809
df_norm
LCL_0811
df_norm
LCL_0813
df_norm
LCL_0821
df_norm
LCL_0823
df_norm
LCL_0824
df_norm
LCL_0826
df_norm
LCL_0827
df_norm
LCL_0828
df_norm
LCL_0831
df_norm
LCL_0834
df_norm
LCL_0835
df_norm
LCL_0839
df_norm
LCL_0847
df_norm
LCL_0848
df_norm
LCL_0853
df_norm
LCL_0854
df_norm
LCL_0858
df_norm
LCL_0863
df_norm
LCL_0865
df_norm
LCL_0870
df_norm
LCL_0874
df_norm
LCL_0879
df_norm
LCL_0880
df_norm
LCL_0883
df_norm

LCL_0903
df_hp
LCL_0905
df_hp
LCL_0907
df_hp
LCL_0911
df_hp
LCL_0912
df_hp
LCL_0916
df_hp
LCL_0923
df_hp
LCL_0935
df_hp
LCL_0937
df_hp
LCL_0942
df_hp
LCL_0950
df_hp
LCL_0958
df_hp
LCL_0959
df_hp
LCL_0960
df_hp
LCL_0961
df_hp
LCL_0963
df_hp
LCL_0967
df_hp
LCL_0975
df_hp
LCL_0978
df_hp
LCL_0980
df_hp
LCL_0984
df_hp
LCL_0987
df_hp
LCL_0989
df_hp
LCL_0999
df_hp
LCL_1002
df_hp
LCL_1006
df_hp
LCL_1007
df_hp
LCL_1025
df_hp
LCL_1027
df_hp
LCL_1028
df_hp
LCL_1036
df_hp
LCL_1041
df_hp
LCL_1048
df_hp
LCL_1055
df_hp
LCL_1066
df_hp
LCL_1069
df_hp
LCL_1078
df_hp
LCL_1086
df_hp

LCL_0884
df_norm
LCL_0887
df_norm
LCL_0888
df_norm
LCL_0889
df_norm
LCL_0890
df_norm
LCL_0891
df_norm
LCL_0898
df_norm
LCL_0904
df_norm
LCL_0909
df_norm
LCL_0914
df_norm
LCL_0917
df_norm
LCL_0920
df_norm
LCL_0921
df_norm
LCL_0922
df_norm
LCL_0926
df_norm
LCL_0928
df_norm
LCL_0943
df_norm
LCL_0954
df_norm
LCL_0964
df_norm
LCL_0968
df_norm
LCL_0969
df_norm
LCL_0982
df_norm
LCL_0990
df_norm
LCL_0994
df_norm
LCL_1005
df_norm
LCL_1021
df_norm
LCL_1032
df_norm
LCL_1034
df_norm
LCL_1035
df_norm
LCL_1039
df_norm
LCL_1043
df_norm
LCL_1050
df_norm
LCL_1076
df_norm
LCL_1077
df_norm
LCL_1079
df_norm
LCL_1087
df_norm
LCL_1088
df_norm
LCL_1097
df_norm

LCL_1090
df_hp
LCL_1094
df_hp
LCL_1096
df_hp
LCL_1098
df_hp
LCL_1100
df_hp
LCL_1102
df_hp
LCL_1104
df_hp
LCL_1105
df_hp
LCL_1109
df_hp
LCL_1110
df_hp
LCL_1115
df_hp
LCL_1119
df_hp
LCL_1120
df_hp
LCL_1121
df_hp
LCL_1123
df_hp
LCL_1124
df_hp
LCL_1125
df_hp
LCL_1128
df_hp
LCL_1133
df_hp
LCL_1134
df_hp
LCL_1135
df_hp
LCL_1137
df_hp
LCL_1142
df_hp
LCL_1143
df_hp
LCL_1145
df_hp
LCL_1147
df_hp
LCL_1152
df_hp
LCL_1154
df_hp
LCL_1156
df_hp
LCL_1159
df_hp
LCL_1163
df_hp
LCL_1164
df_hp
LCL_1166
df_hp
LCL_1172
df_hp
LCL_1173
df_hp
LCL_1175
df_hp
LCL_1177
df_hp
LCL_1182
df_hp
LCL_1188

LCL_1101
df_norm
LCL_1122
df_norm
LCL_1144
df_norm
LCL_1162
df_norm
LCL_1176
df_norm
LCL_1185
df_norm
LCL_1193
df_norm
LCL_1194
df_norm
LCL_1202
df_norm




Figure 1. Annotation diagram of experimental data used in this study. With the colored boxes we show those sub-categories of the data that are compared in our analysis.

3. Results

We have compared the following conditions: Hypertension *versus* Control.

3.1. Identification of target genes

In the first step of the analysis **target genes** were identified from the uploaded experimental data. The most highly methylated genes were used as target genes.

Table 2. Top ten highly methylated genes in Hypertension vs. Control.

[See full table](#) →

ID	Gene description	Gene symbol	Gene schematic representation	Number of methylation sites	Methylation sites in exons	Methylation sites in 5' region
ENSG00000204956	protocadherin gamma subfamily A, 1	PCDHGA1		41	5	0
ENSG00000250349	novel proline rich Gla (G-carboxyglutamic acid) 1 (PRRG1) and tetraspanin 7 (TSPAN7) protein	ENSG00000250349		38	2	2
ENSG00000081853	protocadherin gamma subfamily A, 2	PCDHGA2		37	5	0
ENSG00000204970	protocadherin alpha 1	PCDHA1		34	0	0
ENSG00000254245	protocadherin gamma subfamily A, 3	PCDHGA3		34	5	0
ENSG00000204969	protocadherin alpha 2	PCDHA2		33	3	0
ENSG00000254221	protocadherin gamma subfamily B, 1	PCDHGB1		31	5	0
ENSG00000255408	protocadherin alpha 3	PCDHA3		30	2	0
ENSG00000204967	protocadherin alpha 4	PCDHA4		28	1	0
ENSG00000204965	protocadherin alpha 5	PCDHA5		27	3	1

3.2. Functional classification of genes

A functional analysis of highly methylated genes was done by mapping the genes to several known ontologies, such as Gene Ontology (GO), disease ontology (based on HumanPSD™ database) and the ontology of signal transduction and metabolic pathways from the




Figure 3. Enriched TRANSPATH® Pathways (2023.2) of highly methylated genes in Hypertension vs. Control.

[Full classification →](#)

HumanPSD(TM) disease (2023.2)





Figure 4. Enriched HumanPSD(TM) disease (2023.2) of highly methylated genes in Hypertension vs. Control. The size of the bars correspond to the number of biomarkers of the given disease found among the input set.

[Full classification →](#)

The result of overall Gene Ontology (GO) analysis of the highly methylated genes of the studied pathology can be summarized by the following diagram, revealing the most significant functional categories overrepresented among the observed (highly methylated genes):



3.3. Analysis of enriched transcription factor binding sites and composite modules

In the next step a search for transcription factors binding sites (TFBS) was performed in the regulatory regions of the **target genes** by using the TF binding motif library of the **TRANSFAC®** database. We searched for so called **composite modules** that act as potential condition-specific **enhancers** of the **target genes** in their upstream regulatory regions (-1000 bp upstream of transcription start site (TSS)) and identify transcription factors regulating activity of the genes through such **enhancers**.

Classically, **enhancers** are defined as regions in the genome that increase transcription of one or several genes when inserted in either orientation at various distances upstream or downstream of the gene [8]. Enhancers typically have a length of several hundreds of nucleotides and are bound by multiple transcription factors in a cooperative manner [9].

In the current work we use the Epigenomics data from the track(s) "Methylation track" to predict positions of potential *enhancers* regulating the highly methylated genes revealed by comparative epigenomics analysis. We took genomic regions -550bp upstream and 550bp downstream from the middle point of each interval of the track and check if these regions are located inside the 5kb flanking areas of the highly methylated genes (or inside the body of the genes). In such cases, these genomic regions are used for the search for potential condition-specific enhancers. In all other cases when the differentially expressed genes did not contain epigenomic peaks in their body or in the 5kb flanking regions we used the upstream regulatory regions of these genes (-1000bp upstream and 100bp downstream of TSS) for the search for condition-specific enhancers.

We applied the Composite Module Analyst (CMA) [8] method to detect such potential enhancers, as targets of multiple TFs bound in a cooperative manner to the regulatory regions of the genes of interest. CMA applies a genetic algorithm to construct a generalized model of the enhancers by specifying combinations of TF motifs (from TRANSFAC®) whose sites are most frequently clustered together in the regulatory regions of the studied genes. CMA identifies the transcription factors that through their cooperation provide a synergistic effect and thus have a great influence on the gene regulation process.

Enhancer model potentially involved in regulation of target genes (highly methylated genes in Hypertension vs. Control).

To build the most specific composite modules we choose top methylated genes as the input of CMA algorithm. The obtained CMA model is then applied to compute CMA score for all highly methylated genes in Hypertension vs. Control.

The model consists of 2 module(s). Below, for each module the following information is shown:

- PWMs producing matches,
- number of individual matches for each PWM,
- score of the best match.

Module 1:

V\$GCM1_08 0.92; N=3	V\$ZNF462_01 0.91; N=3	V\$ZNF768_05 0.66; N=3	V\$MZF1_01 1.00; N=2
Module width: 138			

Module 2:

V\$ZBTB33_10 0.99; N=3	V\$ZBTB7B_06 0.88; N=3	V\$NF1C_02 0.94; N=3	V\$BCL6_10 0.73; N=3
Module width: 134			

Model score (-p*log10(pval)): 30.55

Wilcoxon p-value (pval): 9.67e-58

Penalty (p): 0.536

Average yes-set score: 10.31

Average no-set score: 8.60

AUC: 0.74

Separation point: 9.47

False-positive: 31.33%

False-negative: 31.93%

The AUC of the model achieves value significantly higher than expected for a random set of regulatory regions

Z-score = 3.06




Table 3. List of top ten highly methylated genes in Hypertension vs. Control with identified enhancers in their regulatory regions. **CMA score** - the score of the CMA model of the enhancer identified in the regulatory region.

[See full table](#) →

Ensembl IDs	Gene symbol	Gene description	CMA score	Factor names
ENSG00000129965	INS-IGF2	INS-IGF2 readthrough	16.72	BCL-6(h), NF-1C(h), ZBTB7B(h), ZBTB33(h), MZF1(h), GCMA(h), ZNF768(h)...
ENSG00000142303	ADAMTS10	ADAM metallopeptidase with thrombospondin type 1 motif 10	16.24	ZBTB7B(h), GCMA(h), BCL-6(h), ZNF768(h), ZNF462(h), ZBTB33(h), MZF1(h)...
ENSG00000113504	SLC12A7	solute carrier family 12 member 7	16.15	BCL-6(h), ZBTB7B(h), ZNF768(h), GCMA(h), ZNF462(h), ZBTB33(h), MZF1(h)...
ENSG00000251652		novel transcript	15.95	NF-1C(h), ZBTB7B(h), ZNF768(h), ZNF462(h), ZBTB33(h), GCMA(h), MZF1(h)...
ENSG00000284987		novel transcript	15.69	BCL-6(h), ZBTB7B(h), ZBTB33(h), NF-1C(h), ZNF462(h), MZF1(h), ZNF768(h)...
ENSG00000166592	RRAD	RRAD, Ras related glycolysis inhibitor and calcium channel regulator	15.49	NF-1C(h), ZBTB33(h), ZBTB7B(h), GCMA(h), BCL-6(h), ZNF768(h), ZNF462(h)
ENSG00000124212	PTGIS	prostaglandin I2 synthase	15.42	ZBTB33(h), NF-1C(h), ZBTB7B(h), GCMA(h), ZNF768(h), ZNF462(h), BCL-6(h)
ENSG00000171798	KNDC1	kinase non-catalytic C-lobe domain containing 1	15.16	ZBTB7B(h), ZBTB33(h), NF-1C(h), MZF1(h), ZNF768(h), ZNF462(h), GCMA(h)
ENSG00000116981	NT5C1A	5'-nucleotidase, cytosolic IA	15.12	ZNF768(h), ZNF462(h), GCMA(h), ZBTB33(h), ZBTB7B(h), NF-1C(h), MZF1(h)...
ENSG00000269275		novel transcript, antisense to TMEM190	15.06	ZNF768(h), MZF1(h), ZNF462(h), GCMA(h), ZBTB33(h), BCL-6(h), ZBTB7B(h)...


On the basis of the enhancer models we identified transcription factors potentially regulating the **target genes** of our interest. We found 8 transcription factors controlling expression of highly methylated genes in Hypertension vs. Control (see Table 4).

Table 4. Transcription factors of the predicted enhancer model potentially regulating the highly methylated genes (highly methylated genes in Hypertension vs. Control). **Yes-No ratio** is the ratio between frequencies of the sites in Yes sequences versus No sequences. It describes the level of the enrichment of binding sites for the indicated TF in the regulatory target regions. **Regulatory score** is the measure of involvement of the given TF in the controlling of expression of genes that encode master regulators presented below (through positive feedback loops).

[See full table](#) →

ID	Gene symbol	Gene description	Regulatory score	Yes-No ratio
MO000026319	BCL6	BCL6 transcription repressor	2.39	2.82
MO000095587	ZBTB7B	zinc finger and BTB domain containing 7B	2	1.45
MO000092587	ZNF462	zinc finger protein 462	1.75	1.27
MO000026306	GCM1	glial cells missing transcription factor 1	1.47	2.98
MO000092737	ZNF768	zinc finger protein 768	1.26	1.18
MO000024750	NFIC	nuclear factor I C	0	1.45
MO000056592	ZBTB33	zinc finger and BTB domain containing 33	0	1.5
MO000084468	MZF1	myeloid zinc finger 1	0	1.54

The following diagram represents the key transcription factors, which were predicted to be potentially regulating highly methylated genes in the analyzed pathology: BCL6, ZBTB7B and ZNF462.



3.4. Finding master regulators in networks

In the second step of the upstream analysis common regulators of the revealed TFs were identified. These master regulators appear to be the key candidates for therapeutic targets as they have a master effect on regulation of intracellular pathways that activate the pathological process of our study. The identified master regulators are shown in Table 5.

Table 5. Master regulators that may govern the regulation of highly methylated genes in Hypertension vs. Control. **Total rank** is the sum of the ranks of the master molecules sorted by keynode score, CMA score, epigenomics data.

[See full table](#) →

ID	Master molecule name	Gene symbol	Gene description	Total rank
MO000209894	FAM123B-isoform1(h)	AMER1	APC membrane recruitment protein 1	5
MO000209896	FAM123B-isoform2(h)	AMER1	APC membrane recruitment protein 1	6
MO000209895	FAM123B(h)	AMER1	APC membrane recruitment protein 1	7
MO000058267	RSK(h){p}	RPS6KA1, RPS6KA2, RPS6KA3	ribosomal protein S6 kinase A1, ribosomal protein S6 kinase A2, ribosomal protein S6 kinase A3	43
MO000032712	MKP-4(h)	DUSP9	dual specificity phosphatase 9	45
MO000097095	AR-isoform1(h){aceK630}{aceK632}{aceK633}	AR	androgen receptor	59
MO000021454	AR(h)	AR	androgen receptor	60
MO000058849	AR-isoform1(h)	AR	androgen receptor	60
MO000078001	AR(h){sumo}	AR	androgen receptor	60
MO000078150	AR(h){ub}	AR	androgen receptor	60

The intracellular regulatory pathways controlled by the above-mentioned master regulators are depicted in Figure 5. This diagram displays the connections between identified transcription factors, which play important roles in the regulation of highly methylated genes, and selected master regulators, which are responsible for the regulation of these TFs.




Figure 5. Diagram of intracellular regulatory signal transduction pathways of highly methylated genes in Hypertension vs. Control. Master regulators are indicated by red rectangles, transcription factors are blue rectangles, and green rectangles are intermediate molecules, which have been added to the network during the search for master regulators from selected TFs.

[See full diagram →](#)

4. Finding prospective drug targets

The identified master regulators that may govern pathology associated genes were checked for druggability potential using HumanPSD™ [5] database of gene-disease-drug assignments and PASS [11-13] software for prediction of biological activities of chemical compounds on the basis of a (Q)SAR approach. Respectively, for each master regulator protein we have computed two Druggability scores: HumanPSD Druggability score and PASS Druggability score. Where Druggability score represents the number of drugs that are potentially suitable for inhibition (or activation) of the corresponding target either according to the information extracted from medical literature (from HumanPSD™ database) or according to cheminformatics predictions of compounds activity against the examined target (from PASS software).

The cheminformatics druggability check is done using a pre-computed database of spectra of biological activities of chemical compounds from a library of all small molecular drugs from HumanPSD™ database, 2507 pharmaceutically active known chemical compounds in total. The spectra of biological activities has been computed using the program PASS [11-13] on the basis of a (Q)SAR approach.

If both Druggability scores were below defined thresholds (see Methods section for the details) such master regulator proteins were not used in further analysis of drug prediction.

As a result we created the following two tables of prospective drug targets (top targets are shown here):



Table 6. Prospective drug targets selected from full list of identified master regulators filtered by Druggability score from *HumanPSD™* database. **Druggability score** contains the number of drugs that are potentially suitable for inhibition (or activation) of the target. The drug targets are sorted according to the **Total rank** which is the sum of three ranks computed on the basis of the three scores: keynode score, CMA score and expression change score (logFC, if present). See Methods section for details.

[See full table](#) →

Gene symbol	Gene Description	Druggability score	Total rank
AR	androgen receptor	78	60
NTRK3	neurotrophic receptor tyrosine kinase 3	34	155
EPHA8	EPH receptor A8	29	173
TSC22D3	TSC22 domain family member 3	3	241
TSPYL2	TSPY like 2	1	263
ALK	ALK receptor tyrosine kinase	42	275



Table 7. Prospective drug targets selected from full list of identified master regulators filtered by Druggability score predicted by *PASS* software. Here, the **Druggability score** for master regulator proteins is computed as a sum of *PASS* calculated probabilities to be active as a target for various small molecular compounds. The drug targets are sorted according to the **Total rank** which is the sum of three ranks computed on the basis of the three scores: keynode score, CMA score and expression change score (logFC, if present). See Methods section for details.


[See full table](#) →

Gene symbol	Gene Description	Druggability score	Total rank
DUSP9	dual specificity phosphatase 9	35.02	45
AR	androgen receptor	11.8	60
NTRK3	neurotrophic receptor tyrosine kinase 3	9.41	155
EPHA8	EPH receptor A8	17.52	173
ALK	ALK receptor tyrosine kinase	10.66	275
NTF3	neurotrophin 3	17.39	277

Below we represent schematically the main mechanism of the studied pathology. In the schema we considered the top two drug targets of each of the two categories computed above. In addition we have added two top identified master regulators for which no drugs may be identified yet, but that are playing the crucial role in the molecular mechanism of the studied pathology. Thus the molecular mechanism of the studied pathology was predicted to be mainly based on the following key master regulators:

- RSK
- FAM123B
- trkC
- MKP-4
- AR-isoform1

This result allows us to suggest the following schema of affecting the molecular mechanism of the studied pathology:



Drugs which are shown on this schema: estradiol benzoate, 2,5,7-Trihydroxynaphthoquinone, Imatinib, Delta1-dihydrotestosterone, Flavopiridol, Darodipine, Estrone, Progesterone, histamine dihydrochloride and 3-Methyladenine, should be considered as a prospective research initiative for further drug repurposing and drug development. These drugs were selected as top matching treatments to the most prospective drug targets of the studied pathology, however, these results should be considered with special caution and are to be used for research purposes only, as there is not enough clinical information for adapting these results towards immediate treatment of patients.

The drugs given in dark red color on the schema are FDA approved drugs or drugs which have gone through various phases of clinical trials as active treatments against the selected targets.

The drugs given in pink color on the schema are drugs, which were cheminformatically predicted to be active against the selected targets.

5. Identification of potential drugs

In the last step of the analysis we strived to identify known activities as well as drugs with cheminformatically predicted activities that are potentially suitable for inhibition (or activation) of the identified molecular targets in the context of specified human diseases(s).

Proposed drugs are top ranked drug candidates, that were found to be active on the identified targets and were selected from 4 categories:

1. FDA approved drugs or used in clinical trials drugs for the studied pathology;
2. Repurposing drugs used in clinical trials for other pathologies;
3. Drugs, predicted by PASS to be active against identified drug targets and against the studied pathology;
4. Drugs, predicted by PASS to be active against identified drug targets but for other pathologies.

Proposed drugs were selected on the basis of Drug rank which was computed from the ranks sum based on the individual ranks of the following scores:

- Target activity score (depends on ranks of all targets that were found for the selected drug);
- Disease activity score (weighted sum of number of clinical trials on disease(s) under study where the selected drug is known to be applied or PASS Disease activity score - cheminformatically predicted property of the compound to be active against the studied disease(s));
- Clinical validity score (applicable only for drugs predicted on the basis of literature curation in HumanPSD™ database (Tables 8 and 9), reflects the number of the highest clinical trials phase on which the drug was tested for any pathology).

You can refer to the Methods section for more details on drug ranking procedure.

Based on the Drug rank, a numerical value of Drug score was calculated, which reflects the potential activity of the respective drug on the overall molecular mechanism of the studied pathology. Drug score values belong to the range from 1 to 100 and are calculated as a quotient of maximum drug rank and the drug rank of the given drug multiplied by 100.

Top drugs of each category are given in the tables below:

Drugs approved in clinical trials



Table 8. FDA approved drugs or drugs used in clinical trials for the studied pathology (most promising treatment candidates selected for the identified drug targets on the basis of literature curation in [HumanPSD™](#) database)

[See full table](#) →

Name	Target names	Drug score	Disease activity score	Disease trial phase
Imatinib	MAPK10, RPS6KA3, BMPR1A, MARK3, NTRK1, NTRK3, PRKAA2, KDR, FLT4, INSR, LTK, EGFR, ACVR2A, PRKG1, ERBB2, CAMK2B, PRKD1, PKMYT1, IGF1R, BMX, LCK, ALK, PIM2, EPHB2, RPS6KA2, MAPK4, CSNK1G2, PAK3, INSR, BTK, EPHA6, MAPK11, TGFB2, EPHA8, FYN, RET, ABL2, STK3	93	5	Phase 3: Hypertension, Astrocytoma, Bone Marrow Diseases, COVID-19, Familial Primary Pulmonary Hypertension, Fibrosis, Gastrointestinal Stromal Tumors, Glioblastoma, Hematologic Diseases, Hypertension, Pulmonary, Idiopathic Pulmonary Fibrosis, Leukemia, Leukemia, Lymphoid, Leukemia, Myelogenous, Chronic, BCR-ABL Positive, Leukemia, Myeloid, Leukemia, Myeloid, Chronic-Phase, Lung Diseases, Lymphoma, Mucositis, Neoplasms, Precursor Cell Lymphoblastic Leukemia-Lymphoma, Pulmonary Arterial Hypertension, Pulmonary Fibrosis, Recurrence, Sarcoma, Severe Acute Respiratory Syndrome
Pazopanib	MAPK10, RPS6KA3, BMPR1A, MARK3, NTRK1, NTRK3, PRKAA2, KDR, FLT4, INSR, LTK, EGFR, ACVR2A, PRKG1, ERBB2, CAMK2B, PRKD1, PKMYT1, IGF1R, BMX, LCK, ALK, PIM2, EPHB2, RPS6KA2, MAPK4, CSNK1G2, PAK3, INSR, BTK, EPHA6, MAPK11, TGFB2, EPHA8, FYN, RET, ABL2, STK3	91	2	Phase 2: Hypertension, Anemia, Brain Abscess, Brain Neoplasms, Carcinoid Tumor, Carcinoma, Carcinoma, Islet Cell, Carcinoma, Neuroendocrine, Carcinoma, Non-Small-Cell Lung, Carcinoma, Ovarian Epithelial, Carcinoma, Renal Cell, Carcinoma, Small Cell, Carcinoma, Transitional Cell, Central Nervous System Neoplasms, Cholangiocarcinoma, Chondrosarcoma, Chondrosarcoma, Mesenchymal, Corneal Neovascularization, Desmoplastic Small Round Cell Tumor, Dilatation, Pathologic, Drug-Related Side Effects and Adverse Reactions, Edema, Endocrine Gland Neoplasms, Epistaxis, Fibrosarcoma, Gallbladder Neoplasms, Gastrinoma, Gastrointestinal Neoplasms, Gastrointestinal Stromal Tumors, Glioblastoma, Glioma, Gliosarcoma, Glomus Tumor, Glucagonoma, Granular Cell Tumor, Hemangioendothelioma, Hemangioendothelioma, Epithelioid, Hemangiopericytoma, Hemangiosarcoma, Hemorrhage, Histiocytoma, Histiocytoma, Benign Fibrous, Histiocytoma, Malignant Fibrous, Insulinoma, Intestinal Neoplasms, Leiomyosarcoma, Liposarcoma, Lung Neoplasms, Lymphedema, Macular Degeneration, Malignant Carcinoid Syndrome, Melanoma, Multiple Endocrine Neoplasia, Multiple Endocrine Neoplasia Type 1, Multiple Myeloma, Myosarcoma, Neoplasms, Neovascularization, Pathologic, Nerve Sheath Neoplasms, Nervous System Neoplasms, Neuroendocrine Tumors, Neurofibrosarcoma, Osteosarcoma, Ovarian Neoplasms, Pancreatic Neoplasms, Paraganglioma, Paraganglioma, Extra-Adrenal, Pheochromocytoma, Ranula, Recurrence, Rhabdomyosarcoma, Sarcoma, Sarcoma, Alveolar Soft Part, Sarcoma, Clear Cell, Sarcoma, Synovial, Small Cell Lung Carcinoma, Solitary Fibrous Tumors, Somatostatinoma, Stomach Neoplasms, Telangiectasia, Hereditary Hemorrhagic, Telangiectasis, Urinary Bladder Neoplasms
estradiol benzoate	BDNF, AR	87	5	Phase 3: Hypertension, Alzheimer Disease, Anorexia, Anorexia Nervosa, Atrophic Vaginitis, Atrophy, Endometriosis, Essential Hypertension, Genetic Diseases, Inborn, Gonadal Dysgenesis, Hemorrhage, Hot Flashes, Hypogonadism, Infertility, Infertility, Female, Menopause, Premature, Metrorrhagia, Premature Birth, Primary Ovarian Insufficiency, Syndrome, Turner Syndrome, Uterine Hemorrhage, Vaginitis
Drospirenone	AR	87	5	Phase 3: Hypertension, Acne Vulgaris, Cysts, Dysbiosis, Essential Hypertension, Hot Flashes, Neural Tube Defects, Polycystic Ovary Syndrome, Spinal Dysraphism, Syndrome
Sorafenib	MAPK10, RPS6KA3, BMPR1A, MARK3, NTRK1, NTRK3, PRKAA2, KDR, FLT4, INSR, LTK, EGFR, ACVR2A, PRKG1, ERBB2, CAMK2B, PRKD1, PKMYT1, IGF1R, BMX, LCK, ALK, PIM2, EPHB2, RPS6KA2, MAPK4,	87	1	Phase 1: Hypertension, Adenocarcinoma, Adenoma, Adenoma, Liver Cell, Astrocytoma, Bile Duct Neoplasms, Biliary Tract Neoplasms, Blast Crisis, Brain Abscess, Brain Neoplasms, Breast Neoplasms, Carcinoid Tumor, Carcinoma, Carcinoma, Hepatocellular, Carcinoma, Non-Small-Cell Lung, Carcinoma, Renal Cell, Cholangiocarcinoma, Familial Primary Pulmonary Hypertension, Gallbladder Neoplasms, Glioblastoma, Glioma, Gliosarcoma, Hate, Hematologic Neoplasms, Hodgkin Disease, Hypertension, Pulmonary, Hypophosphatemia, Kidney Diseases, Kidney Neoplasms, Leukemia, Leukemia, Lymphocytic, Chronic, B-Cell, Leukemia, Lymphoid, Leukemia, Myelogenous, Chronic, BCR-ABL Positive, Leukemia, Myeloid, Leukemia, Myeloid, Accelerated Phase, Leukemia, Myeloid, Acute, Leukemia, Myeloid, Chronic, Atypical, BCR-ABL Negative, Leukemia, Myeloid, Chronic-Phase, Leukemia, Myelomonocytic, Chronic, Leukemia,

CSNK1G2, PAK3,
INSRR, PRKCZ,
BTK, EPHA6,
MAPK11, TGFB2,
EPHA8, FYN, RET,
ABL2, STK3

Myelomonocytic, Juvenile, Leukemia, Promyelocytic, Acute, Liver Neoplasms, Lung Neoplasms, Lymphoma, Lymphoma, Non-Hodgkin, Melanoma, Mesothelioma, Multiple Myeloma, Myelodysplastic Syndromes, Myelodysplastic-Myeloproliferative Diseases, Myeloproliferative Disorders, Myosarcoma, Neoplasms, Neoplasms, Plasma Cell, Neuroblastoma, Neuroendocrine Tumors, Oligodendroglioma, Osteosarcoma, Pancreatic Neoplasms, Pica, Plasmacytoma, Precursor Cell Lymphoblastic Leukemia-Lymphoma, Preleukemia, Primary Myelofibrosis, Pulmonary Arterial Hypertension, Recurrence, Rhabdomyosarcoma, Sarcoma, Sarcoma, Ewing, Syndrome, Thrombocytosis, Urinary Bladder Neoplasms

The **Disease trial phase** column reflects the maximum clinical trials phase in which the drug was studied for the analyzed pathology.

Repurposing drugs



Table 9. Repurposed drugs used in clinical trials for other pathologies (prospective drugs against the identified drug targets on the basis of literature curation in *HumanPSD™* database)

[See full table](#) →

Name	Target names	Drug score	Maximum trial phase
Flavopiridol	MAPK10, RPS6KA3, CDK6, BMPR1A, MARK3, NTRK1, NTRK3, PRKAA2, KDR, FLT4, INSR, LTK, EGFR, ACVR2A, PRKG1, ERBB2, CAMK2B, PRKD1, PKMYT1, IGF1R, BMX, LCK, ALK, PIM2, EPHB2, RPS6KA2, MAPK4, CSNK1G2, PAK3, INSRR, BTK, EPHA6, XIAP, MAPK11, TGFB2, EPHA8, FYN, RET, ABL2, STK3	87	Phase 2: Embolism, Head and Neck Neoplasms, Lymphoma, Lymphoma, B-Cell, Lymphoma, Large B-Cell, Diffuse, Lymphoma, Mantle-Cell, Lymphoma, Non-Hodgkin, Neoplasms, Sarcoma, Thromboembolism
CI-1033	MAPK10, RPS6KA3, BMPR1A, MARK3, NTRK1, NTRK3, PRKAA2, KDR, FLT4, INSR, LTK, EGFR, ACVR2A, PRKG1, ERBB2, CAMK2B, PRKD1, PKMYT1, IGF1R, BMX, LCK, ALK, PIM2, EPHB2, RPS6KA2, MAPK4, CSNK1G2, ERBB4, PAK3, INSRR, BTK, EPHA6, MAPK11, TGFB2, EPHA8, FYN, RET, ABL2, STK3	87	Phase 2: Breast Neoplasms, Carcinoma, Non-Small-Cell Lung, Lung Neoplasms, Neoplasms
pelitinib	MAPK10, RPS6KA3, BMPR1A, MARK3, NTRK1, NTRK3, PRKAA2, KDR, FLT4, INSR, LTK, EGFR, ACVR2A, PRKG1, ERBB2, CAMK2B, PRKD1, PKMYT1, IGF1R, BMX, LCK, ALK, PIM2, EPHB2, RPS6KA2, MAPK4, CSNK1G2, ERBB4, PAK3, INSRR, BTK, EPHA6, MAPK11, TGFB2, EPHA8, FYN, RET, ABL2, STK3	87	Phase 2: Carcinoma, Non-Small-Cell Lung, Colonic Neoplasms, Colorectal Neoplasms, Neoplasms, Rectal Neoplasms
bms-387032	MAPK10, RPS6KA3, BMPR1A, MARK3, NTRK1, NTRK3, PRKAA2, KDR, FLT4, INSR, LTK, EGFR, ACVR2A, PRKG1, ERBB2, CAMK2B, PRKD1, PKMYT1, IGF1R, BMX, LCK, ALK, PIM2, EPHB2, PRKCG, RPS6KA2, MAPK4, CSNK1G2, PAK3, INSRR, BTK, EPHA6, MAPK11, TGFB2, EPHA8, FYN, RET, ABL2, STK3	86	Phase 1: Leukemia, Lymphocytic, Chronic, B-Cell, Lymphoma, Mantle-Cell, Multiple Myeloma, Neoplasms
Dasatinib	MAPK10, RPS6KA3, BMPR1A, MARK3, NTRK1, NTRK3, PRKAA2, KDR, FLT4, INSR, LTK, EGFR, ACVR2A, PRKG1, ERBB2, CAMK2B, PRKD1, PKMYT1, IGF1R, BMX, LCK, ALK, PIM2, EPHB2, RPS6KA2, MAPK4, CSNK1G2, PAK3, INSRR, PRKCZ, BTK, EPHA6, MAPK11, TGFB2, EPHA8, FYN, RET, ABL2, STK3	86	Phase 4: Leukemia, Leukemia, Myelogenous, Chronic, BCR-ABL Positive, Leukemia, Myeloid, Precursor Cell Lymphoblastic Leukemia-Lymphoma

The **Maximum trial phase** column reflects the maximum clinical trials phase in which the drug was studied for any pathology.



Table 10. Prospective drugs, predicted by *PASS* software to be active against the identified drug targets with predicted activity against the studied disease(s) (drug candidates predicted with the cheminformatics tool *PASS*)

[See full table](#) →

Name	Target names	Drug score	Target activity score
Moexipril	BDNF, NTF3, ITGB3	91	0.35
Quinapril	BDNF, NTF3, ITGB3	89	0.27
Nicergoline	PRKAA2, PRKG1, PKMYT1	81	8.58E-2
Hesperetin	MAPK10, BDNF, MAPK4, MAPK11, NTF3	77	7.17E-2
(2s,3s)-Trans-Dihydroquercetin	MAPK10, BDNF, MAPK4, MAPK11, NTF3	75	6.18E-2



Table 11. Prospective drugs, predicted by PASS software to be active against the identified drug targets, though without cheminformatically predicted activity against the studied disease(s) (drug candidates predicted with the cheminformatics tool PASS)
See full table →

Name	Target names	Drug score	Target activity score
2,5,7-Trihydroxynaphthoquinone	MAPK10, DUSP22, MAPK4, CDC14B, MAPK11, DUSP14, DUPD1, DUSP9, BRCA1	86	0.25
3-METHYL-1,6,8-TRIHYDROXYANTHRAQUINONE	MAPK10, DUSP22, MAPK4, CDC14B, MAPK11, DUSP14, DUPD1, DUSP9, BRCA1	86	0.21
[[N-(Benzyloxycarbonyl)Amino]Methyl]Phosphate	PTPRR, ACE2, GRIN1, DUSP22, PTPRO, PTPRF, PTPN2, PTPN13, PTPRH, CDC14B, DUSP14, DUPD1, DUSP9, PTPN21	84	0.49
histamine dihydrochloride	BMX, LCK, MAPK10, ALK, EPHB2, NTRK1, NTRK3, MAPK4, KDR, ERBB4, INSR, FLT4, INSR, LTK, EGFR, BTK, EPHA6, MAPK11, ERBB2, FYN, EPHA8, RET, ABL2, IGF1R	84	1.21
Fenoldopam	MAPK10, GRIN1, MAPK4, MAPK11	82	0.14

As the result of drug search we propose the following drugs as most promising candidates for treating the pathology under study: Imatinib, Flavopiridol, Moexipril and 2,5,7-Trihydroxynaphthoquinone. These drugs were selected for acting on the following targets: NTRK3, NTF3 and DUSP9, which were predicted to be active in the molecular mechanism of the studied pathology.

The selected drugs are top ranked drug candidates from each of the four categories of drugs: (1) FDA approved drugs or used in clinical trials drugs for the studied pathology; (2) repurposing drugs used in clinical trials for other pathologies; (3) drugs, predicted by PASS software to be active against the studied pathology; (4) drugs, predicted by PASS software to be repurposed from other pathologies.

6. Conclusion

We applied the software package "Genome Enhancer" to a data set that contains *epigenomics* data. The study is done in the context of *Hypertension*. The data were pre-processed, statistically analyzed and highly methylated genes were identified. Also checked was the enrichment of GO or disease categories among the studied gene sets.

We propose the following drugs as most promising candidates for treating the pathology under study:

Imatinib, Flavopiridol, Moexipril and 2,5,7-Trihydroxynaphthoquinone

These drugs were selected for acting on the following targets: NTRK3, NTF3 and DUSP9, which were predicted to be involved in the molecular mechanism of the pathology under study.

The identified molecular mechanism of the studied pathology was predicted to be mainly based on the following key drug targets:

RSK, FAM123B, trkC, MKP-4 and AR-isoform1

These potential drug targets should be considered as a prospective research initiative for further drug repurposing and drug development purposes. The following drugs were predicted as, matching those drug targets: estradiol benzoate, 2,5,7-Trihydroxynaphthoquinone, Imatinib, Delta1-dihydrotestosterone, Flavopiridol, Darodipine, Estrone, Progesterone, histamine dihydrochloride and 3-Methyladenine. These drugs should be considered with special caution for research purposes only.

In this study, we came up with a detailed signal transduction network regulating highly methylated genes in the studied pathology. In this network we have revealed the following top master regulators (signaling proteins and their complexes) that play a crucial role in the molecular mechanism of the studied pathology, which can be proposed as the most promising molecular targets for further drug repurposing and drug development initiatives.

- RSK
- FAM123B
- trkC
- MKP-4
- AR-isoform1

Potential drug compounds which can be affecting these targets can be found in the "Finding prospective drug targets" section.

7. Methods

Databases used in the study

Transcription factor binding sites in promoters and enhancers of highly methylated genes were analyzed using known DNA-binding motifs described in the [TRANSFAC®](https://genexplain.com/transfac) library, release 2023.2 (geneXplain GmbH, Wolfenbüttel, Germany) (<https://genexplain.com/transfac>).

The master regulator search uses the [TRANSPATH®](https://genexplain.com/transpath) database (BIOBASE), release 2023.2 (geneXplain GmbH, Wolfenbüttel, Germany) (<https://genexplain.com/transpath>). A comprehensive signal transduction network of human cells is built by the software on the basis of reactions annotated in [TRANSPATH®](https://genexplain.com/transpath).

The information about drugs corresponding to identified drug targets and clinical trials references were extracted from [HumanPSD™](https://genexplain.com/humanpsd) database, release 2023.2 (<https://genexplain.com/humanpsd>).

The Ensembl database release Human104.38 (hg38) (<http://www.ensembl.org>) was used for gene IDs representation and Gene Ontology (GO) (<http://geneontology.org>) was used for functional classification of the studied gene set.

Epigenomics data processing

When analyzing a list of CpG sites, we compute the fold change values between the methylation status in the studied pathology and the control set. Top 10 000 CpG sites with highest logFC values are taken to further analysis. These sites are mapped to corresponding genes, which will be further compared to the list of housekeeping genes at the step of promoter analysis.

Methods for the analysis of enriched transcription factor binding sites and composite modules

Transcription factor binding sites in promoters and enhancers of differentially expressed genes were analyzed using known DNA-binding motifs. The motifs are specified using position weight matrices (PWMs) that give weights to each nucleotide in each position of the DNA binding motif for a transcription factor or a group of them.

We search for transcription factor binding sites (TFBS) that are enriched in the enhancers under study as compared to a background set of promoters of housekeeping genes. We denote study and background sets briefly as Yes and No sets. In the current work we used a workflow considering promoter sequences of a standard length of 1100 bp (-1000 to +100). The error rate in this part of the pipeline is controlled by estimating the adjusted p-value (using the Benjamini-Hochberg procedure) in comparison to the TFBS frequency found in randomly selected regions of the human genome (adj.p-value < 0.01).

We have applied the CMA algorithm (Composite Module Analyst) for searching composite modules [7] in the promoters and enhancers of the Yes and No sets. We searched for a composite module consisting of a cluster of 10 TFs in a sliding window of 200-300 bp that statistically significantly separates sequences in the Yes and No sets (minimizing Wilcoxon p-value).

Methods for finding master regulators in networks

We searched for master regulator molecules in signal transduction pathways upstream of the identified transcription factors. The master regulator search uses a comprehensive signal transduction network of human cells. The main algorithm of the master regulator search has been described earlier [3,4]. The goal of the algorithm is to find nodes in the global signal transduction network that may potentially regulate the activity of a set of transcription factors found at the previous step of the analysis. Such nodes are considered as most promising drug targets, since any influence on such a node may switch the transcriptional programs of hundreds of genes that are regulated by the respective TFs. In our analysis, we have run the algorithm with a maximum radius of 12 steps upstream of each TF in the input set. The error rate of this algorithm is controlled by applying it 10000 times to randomly generated sets of input transcription factors of the same set-size. Z-score and FDR value of ranks are calculated then for each potential master regulator node on the basis of such random runs (see detailed description in [9]). We control the error rate by the FDR threshold 0.05.

Methods for analysis of pharmaceutical compounds

We seek for the optimal combination of molecular targets (key elements of the regulatory network of the cell) that potentially interact with pharmaceutical compounds from a library of known drugs and biologically active chemical compounds, using information about known drugs from HumanPSD™ and predicting potential drugs using [PASS](#) program.

Method for analysis of known pharmaceutical compounds

We selected compounds from HumanPSD™ database that have at least one target. Next, we sort compounds using "Drug rank" that is the sum of the following ranks:

1. ranking by "Target activity score" ($T\text{-score}_{PSD}$),
2. ranking by "Disease activity score" ($D\text{-score}_{PSD}$),
3. ranking by "Clinical validity score".

"Target activity score" ($T\text{-score}_{PSD}$) is calculated as follows:

$$T\text{-score}_{PSD} = -\frac{|T|}{|T| + w(|AT| - |T|)} \sum_{t \in T} \log_{10} \left(\frac{\text{rank}(t)}{1 + \max\text{Rank}(T)} \right),$$

where T is set of all targets related to the compound intersected with input list, $|T|$ is number of elements in T , AT and $|AT|$ are set set of all targets related to the compound and number of elements in it, w is weight multiplier, $\text{rank}(t)$ is rank of given target, $\max\text{Rank}(T)$ equals $\max(\text{rank}(t))$ for all targets t in T .

We use following formula to calculate "Disease activity score" ($D\text{-score}_{PSD}$):

$$D\text{-score}_{PSD} = \begin{cases} \sum_{d \in D} \sum_{p \in P} \text{phase}(d, p) \\ 0, D = \emptyset \end{cases},$$

where D is the set of selected diseases, and if D is empty set, $D\text{-score}_{PSD}=0$. P is a set of all known phases for each disease, $\text{phase}(p,d)$ equals to the phase number if there are known clinical trials for the selected disease on this phase and zero otherwise.

The clinical validity score reflects the number of the highest clinical trials phase (from 1 to 4) on which the drug was ever tested for any pathology.

Method for prediction of pharmaceutical compounds

In this study, the focus was put on compounds with high pharmacological efficiency and low toxicity. For this purpose, comprehensive library of chemical compounds and drugs was subjected to a SAR/QSAR analysis. This library contains 13040 compounds along with their pre-calculated potential pharmacological activities of those substances, their possible side and toxic effects, as well as the possible mechanisms of action. All biological activities are expressed as probability values for a substance to exert this activity (Pa).

We selected compounds that satisfied the following conditions:

1. Toxicity below a chosen toxicity threshold (defines as Pa , probability to be active as toxic substance).
2. For all predicted pharmacological effects that correspond to a set of user selected disease(s) Pa is greater than a chosen effect threshold.
3. There are at least 2 targets (corresponding to the predicted activity-mechanisms) with predicted Pa greater than a chosen target threshold.

The maximum Pa value for all toxicities corresponding to the given compound is selected as the "Toxicity score". The maximum Pa value for all activities corresponding to the selected diseases for the given compound is used as the "Disease activity score". "Target activity score" ($T\text{-score}$) is calculated as follows:

$$T\text{-score}(s) = \frac{|T|}{|T| + w(|AT| - |T|)} \sum_{m \in M(s)} \left(pa(m) \sum_{g \in G(m)} IAP(g) \text{optWeight}(g) \right),$$

where $M(s)$ is the set of activity-mechanisms for the given structure (which passed the chosen threshold for activity-mechanisms Pa); $G(m)$ is the set of targets (converted to genes) that corresponds to the given activity-mechanism (m) for the given compound; $pa(m)$ is the probability to be active of the activity-mechanism (m), $IAP(g)$ is the invariant accuracy of prediction for gene from $G(m)$; $\text{optWeight}(g)$ is the additional weight multiplier for gene. T is set of all targets related to the compound intersected with input list, $|T|$ is number of elements in T , AT and $|AT|$ are set set of all targets related to the compound and number of elements in it, w is weight multiplier.

"Druggability score" ($D\text{-score}$) is calculated as follows:

$$D\text{-score}(g) = IAP(g) \sum_{s \in S(g)} \sum_{m \in M(s,g)} pa(m),$$

where $S(g)$ is the set of structures for which target list contains given target, $M(s,g)$ is the set of activity-mechanisms (for the given structure) that corresponds to the given gene, $pa(m)$ is the probability to be active of the activity-mechanism (m), $IAP(g)$ is the invariant accuracy of prediction for the given gene.

8. References

1. Kel A, Voss N, Jauregui R, Kel-Margoulis O, Wingender E. Beyond microarrays: Finding key transcription factors controlling signal transduction pathways. *BMC Bioinformatics*. **2006**;7(S2), S13. doi:10.1186/1471-2105-7-s2-s13
2. Stegmaier P, Voss N, Meier T, Kel A, Wingender E, Borlak J. Advanced Computational Biology Methods Identify Molecular Switches for Malignancy in an EGF Mouse Model of Liver Cancer. *PLoS ONE*. **2011**;6(3):e17738. doi:10.1371/journal.pone.0017738
3. Koschmann J, Bhar A, Stegmaier P, Kel A, Wingender E. “Upstream Analysis”: An Integrated Promoter-Pathway Analysis Approach to Causal Interpretation of Microarray Data. *Microarrays*. **2015**;4(2):270-286. doi:10.3390/microarrays4020270.
4. Kel A, Stegmaier P, Valeev T, Koschmann J, Poroikov V, Kel-Margoulis OV, and Wingender E. Multi-omics “upstream analysis” of regulatory genomic regions helps identifying targets against methotrexate resistance of colon cancer. *EuPA Open Proteom*. **2016**;13:1-13. doi:10.1016/j.euprot.2016.09.002
5. Michael H, Hogan J, Kel A et al. Building a knowledge base for systems pathology. *Brief Bioinformatics*. **2008**;9(6):518-531. doi:10.1093/bib/bbn038
6. Matys V, Kel-Margoulis OV, Fricke E, Liebich I, Land S, Barre-Dirrie A, Reuter I, Chekmenev D, Krull M, Hornischer K, Voss N, Stegmaier P, Lewicki-Potapov B, Saxel H, Kel AE, Wingender E. TRANSFAC and its module TRANSCompel: transcriptional gene regulation in eukaryotes. *Nucleic Acids Res*. **2006**;34(90001):D108-D110. doi:10.1093/nar/gkj143
7. Kel AE, Gössling E, Reuter I, Cheremushkin E, Kel-Margoulis OV, Wingender E. MATCH: A tool for searching transcription factor binding sites in DNA sequences. *Nucleic Acids Res*. **2003**;31(13):3576-3579. doi:10.1093/nar/gkg585
8. Waleev T, Shtokalo D, Konovalova T, Voss N, Cheremushkin E, Stegmaier P, Kel-Margoulis O, Wingender E, Kel A. Composite Module Analyst: identification of transcription factor binding site combinations using genetic algorithm. *Nucleic Acids Res*. **2006**;34(Web Server issue):W541-5.
9. Krull M, Pistor S, Voss N, Kel A, Reuter I, Kronenberg D, Michael H, Schwarzer K, Potapov A, Choi C, Kel-Margoulis O, Wingender E. TRANSPATH: an information resource for storing and visualizing signaling pathways and their pathological aberrations. *Nucleic Acids Res*. **2006**;34(90001):D546-D551. doi:10.1093/nar/gkj107
10. Boyarskikh U, Pintus S, Mandrik N, Stelmashenko D, Kiselev I, Evshin I, Sharipov R, Stegmaier P, Kolpakov F, Filipenko M, Kel A. Computational master-regulator search reveals mTOR and PI3K pathways responsible for low sensitivity of NCI-H292 and A427 lung cancer cell lines to cytotoxic action of p53 activator Nutlin-3. *BMC Med Genomics*. **2018**;11(1):12. doi:10.1186/1471-2105-7-s2-s13
11. Filimonov D, Poroikov V. Probabilistic Approaches in Activity Prediction. Varnek A, Tropsha A. *Cheminformatics Approaches to Virtual Screening*. Cambridge (UK): RSC Publishing. **2008**;:182-216.
12. Filimonov DA, Poroikov VV. Prognosis of specters of biological activity of organic molecules. *Russian chemical journal*. **2006**;50(2):66-75 (russ)
13. Filimonov D, Poroikov V, Borodina Y, Glorizova T. Chemical Similarity Assessment Through Multilevel Neighborhoods of Atoms: Definition and Comparison with the Other Descriptors. *ChemInform*. **1999**;39(4):666-670. doi:10.1002/chin.199940210

Thank you for using the Genome Enhancer!

In case of any questions please contact us at support@genexplain.com

Supplementary material

1. [Supplementary table 1 - Detailed report. Composite modules and master regulators \(highly methylated genes in Hypertension vs. Control\).](#)
2. [Supplementary table 2 - Detailed report. Pharmaceutical compounds and drug targets.](#)

Disclaimer

Decisions regarding care and treatment of patients should be fully made by attending doctors. The predicted chemical compounds listed in the report are given only for doctor’s consideration and they cannot be treated as prescribed medication. It is the physician’s responsibility to independently decide whether any, none or all of the predicted compounds can be used solely or in combination for patient treatment purposes, taking into account all applicable information regarding FDA prescribing recommendations for any therapeutic and the patient’s condition, including, but not limited to, the patient’s and family’s medical history, physical examinations, information from various diagnostic tests, and patient preferences in accordance with the current standard of care. Whether or not a particular patient will benefit from a selected therapy is based on many factors and can vary significantly.

The compounds predicted to be active against the identified drug targets in the report are not guaranteed to be active against any particular patient’s condition. GeneXplain GmbH does not give any assurances or guarantees regarding the treatment information and conclusions given in the report. There is no guarantee that any third party will provide a refund for any of the treatment decisions made based on these results. None of the listed compounds was checked by Genome Enhancer for adverse side-effects or even toxic effects.

The analysis report contains information about chemical drug compounds, clinical trials and disease biomarkers retrieved from the HumanPSD™ database of gene-disease assignments maintained and exclusively distributed worldwide by geneXplain GmbH. The information contained in this database is collected from scientific literature and public clinical trials resources. It is updated to the best of geneXplain's knowledge however we do not guarantee completeness and reliability of this information leaving the final checkup and consideration of the predicted therapies to the medical doctor.

The scientific analysis underlying the Genome Enhancer report employs a complex analysis pipeline which uses geneXplain's proprietary Upstream Analysis approach, integrated with TRANSFAC® and TRANSPATH® databases maintained and exclusively distributed worldwide by geneXplain GmbH. The pipeline and the databases are updated to the best of geneXplain's knowledge and belief, however, geneXplain GmbH shall not give a warranty as to the characteristics or to the content and any of the results produced by Genome Enhancer. Moreover, any warranty concerning the completeness, up-to-dateness, correctness and usability of Genome Enhancer information and results produced by it, shall be excluded.

The results produced by Genome Enhancer, including the analysis report, severely depend on the quality of input data used for the analysis. It is the responsibility of Genome Enhancer users to check the input data quality and parameters used for running the Genome Enhancer pipeline.

Note that the text given in the report is not unique and can be fully or partially repeated in other Genome Enhancer analysis reports, including reports of other users. This should be considered when publishing any results or excerpts from the report. This restriction refers only to the general description of analysis methods used for generating the report. All data and graphics referring to the concrete set of input data, including lists of mutated genes, differentially expressed genes/proteins/metabolites, functional classifications, identified transcription factors and master regulators, constructed molecular networks, lists of chemical compounds and reconstructed model of molecular mechanisms of the studied pathology are unique in respect to the used input data set and Genome Enhancer pipeline parameters used for the current run.