

ITGA6 and ITGB6 are promising druggable targets for treating Squamous Cell Carcinoma that control activity of SMAD3, TP63 and GTF2I transcription factors on promoters of differentially expressed genes

Demo User

geneXplain GmbH

info@genexplain.com

Data received on 13/08/2019 ; Run on 26/06/2024 ; Report generated on 27/06/2024

Genome Enhancer release 3.4 (TRANSFAC®, TRANSPATH® and HumanPSD™ release 2024.1)



Abstract

In the present study we applied the software package "Genome Enhancer" to a data set that contains *transcriptomics* data. The study is done in the context of *Squamous Cell Carcinoma*. The goal of this pipeline is to identify potential drug targets in the molecular network that governs the studied pathological process. In the first step of analysis pipeline discovers transcription factors (TFs) that regulate genes activities in the pathological state. The activities of these TFs are controlled by so-called master regulators, which are identified in the second step of analysis. After a subsequent druggability checkup, the most promising master regulators are chosen as potential drug targets for the analyzed pathology. At the end the pipeline comes up with (a) a list of known drugs and (b) investigational active chemical compounds with the potential to interact with selected drug targets.

From the data set analyzed in this study, we found the following TFs to be potentially involved in the regulation of the differentially expressed genes: SMAD3, TP63, TP53, GTF2I, SP1 and JUND. The subsequent network analysis suggested

- integrins
- ornithine decarboxylase
- EGFR
- EGFR

as the most promising molecular targets for further research, drug development and drug repurposing initiatives on the basis of identified molecular mechanism of the studied pathology. Having checked the actual druggability potential of the full list of identified targets, both, via information available in medical literature and via cheminformatics analysis of drug compounds, we have identified the following drugs as the most promising treatment candidates for the studied pathology: Erlotinib, Curcumin and $\{(2Z)-4\text{-AMINO-2-}[(4\text{-METHOXYPHENYL})\text{IMINO}]\text{-2,3-DIHYDRO-1,3-THIAZOL-5-YL}\}(4\text{-METHOXYPHENYL})\text{METHANONE}$.

1. Introduction

Recording "-omics" data to measure gene activities, protein expression or metabolic events is becoming a standard approach to characterize the pathological state of an affected organism or tissue. Increasingly, several of these methods are applied in a combined approach leading to large "multiomics" datasets. Still the challenge remains how to reveal the underlying molecular mechanisms that render a given pathological state different from the norm. The disease-causing mechanism can be described by a re-wiring of the cellular regulatory network, for instance as a result of a genetic or epigenetic alterations influencing the activity of relevant genes. Reconstruction of the disease-specific regulatory networks can help identify potential master regulators of the respective pathological process. Knowledge about these master regulators can point to ways how to block a pathological regulatory cascade. Suppression of certain molecular targets as components of these cascades may stop the pathological process and cure the disease.

Conventional approaches of statistical "-omics" data analysis provide only very limited information about the causes of the observed phenomena and therefore contribute little to the understanding of the pathological molecular mechanism. In contrast, the "upstream analysis" method [1-4] applied here has been devised to provide a casual interpretation of the data obtained for a pathology state. This approach comprises two major steps: (1) analysing promoters and enhancers of differentially expressed genes for the transcription factors (TFs) involved in their regulation and, thus, important for the process under study; (2) re-constructing the signaling pathways that activate these TFs and identifying master regulators at the top of such pathways. For the first step, the database TRANSFAC® [6] is employed together with the TF binding site identification algorithms Match [7] and CMA [8]. The second step involves the signal transduction database TRANSPATH® [9] and special graph search algorithms [10] implemented in the software "Genome Enhancer".

The "upstream analysis" approach has now been extended by a third step that reveals known drugs suitable to inhibit (or activate) the identified molecular targets in the context of the disease under study. This step is performed by using information from HumanPSD™ database [5]. In addition, some known drugs and investigational active chemical compounds are subsequently predicted as potential ligands for the revealed molecular targets. They are predicted using a pre-computed database of spectra of biological activities of chemical compounds of a library of 2245 known drugs and investigational chemical compounds from HumanPSD™ database. The spectra of biological activities for these compounds are computed using the program PASS on the basis of a (Q)SAR approach [11-13]. These predictions can be used for the research purposes - for further drug development and drug repurposing initiatives.

2. Data

For this study the following experimental data was used:

Table 1. Experimental datasets used in the study

File name	Data type
SRR349741.fastq	Transcriptomics
SRR349742.fastq	Transcriptomics
SRR349748.fastq	Transcriptomics
SRR349749.fastq	Transcriptomics




Figure 1. Annotation diagram of experimental data used in this study. With the colored boxes we show those sub-categories of the data that are compared in our analysis.

3. Results

We have compared the following conditions: Experiment: Squamous Cell Carcinoma versus Control: Non-tumour tissue.

3.1. Identification of target genes

In the first step of the analysis **target genes** were identified from the uploaded experimental data. We applied the edgeR tool (R/Bioconductor package integrated into our pipeline) and compared gene expression in the following sets: "Experiment: Squamous Cell Carcinoma" with "Control: Non-tumour tissue". edgeR calculated the LogFC (the logarithm to the base 2 of the fold change between different conditions), the p-value and the adjusted p-value (corrected for multiple testing) of the observed fold change. As a result, we detected 1955 upregulated genes (LogFC>0.1) out of which 768 genes were found as significantly upregulated (p-value<0.1) and 1739 downregulated genes (LogFC<-0.1) out of which 650 genes were significantly downregulated (p-value<0.1). See tables below for the top significantly up- and downregulated genes. Below we call **target genes** the full list of up- and downregulated genes revealed in our analysis (see tables in [Supplementary section](#)).

Table 2. Top ten significant **up-regulated** genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue.

[See full table](#) →

ID	Gene symbol	Gene description	logFC	logCPM	PValue	FDR
ENSG00000115758	ODC1	ornithine decarboxylase 1	6.73	10.37	6.78E-9	6.85E-7
ENSG00000148053	NTRK2	neurotrophic receptor tyrosine kinase 2	5.99	9.33	1.47E-9	1.99E-7
ENSG00000113140	SPARC	secreted protein acidic and cysteine rich	5.74	10.75	1.44E-7	9.8E-6
ENSG00000163359	COL6A3	collagen type VI alpha 3 chain	5.19	9.2	1.54E-5	4.36E-4
ENSG00000120708	TGFBI	transforming growth factor beta induced	4.81	8.83	1.53E-9	2.01E-7
ENSG00000134871	COL4A2	collagen type IV alpha 2 chain	4.69	8.02	9.35E-10	1.36E-7
ENSG00000186340	THBS2	thrombospondin 2	4.67	8.54	6.35E-5	1.34E-3
ENSG00000146648	EGFR	epidermal growth factor receptor	4.44	9.65	3.25E-4	4.84E-3
ENSG00000145824	CXCL14	C-X-C motif chemokine ligand 14	4.43	8.61	2.44E-5	6.33E-4
ENSG00000187134	AKR1C1	aldo-keto reductase family 1 member C1	4.41	9.04	1.06E-10	2.88E-8

Table 3. Top ten significant **down-regulated** genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue. [See full table](#) →

ID	Gene symbol	Gene description	logFC	logCPM	PValue	FDR
ENSG00000136155	SCEL	sciellin	-7.72	11.12	2.73E-15	5.38E-12
ENSG00000163209	SPRR3	small proline rich protein 3	-6.69	14.45	8.44E-4	1.1E-2
ENSG00000143369	ECM1	extracellular matrix protein 1	-6.38	11.04	4.35E-10	7.45E-8
ENSG00000189334	S100A14	S100 calcium binding protein A14	-6.37	10.46	1.1E-10	2.88E-8
ENSG00000229732		novel transcript	-6.27	12.97	4.93E-12	2.42E-9
ENSG00000086548	CEACAM6	CEA cell adhesion molecule 6	-6.2	10.31	5.18E-14	4.37E-11
ENSG00000171401	KRT13	keratin 13	-6.15	14.93	8.06E-11	2.44E-8
ENSG00000087128	TMPRSS11E	transmembrane serine protease 11E	-5.98	10.11	6.26E-9	6.48E-7
ENSG00000197632	SERPINB2	serpin family B member 2	-5.86	8.73	5.56E-14	4.37E-11
ENSG00000165272	AQP3	aquaporin 3 (Gill blood group)	-5.81	11.35	5.75E-5	1.23E-3


3.2. Functional classification of genes

A functional analysis of differentially expressed genes was done by mapping the significant up-regulated and significant down-regulated genes to several known ontologies, such as Gene Ontology (GO), disease ontology (based on HumanPSD™ database) and the ontology of signal transduction and metabolic pathways from the [TRANSPATH®](#) database. Statistical significance was computed using a binomial test.


Figures 3-8 show the most significant categories.

Heatmap of differentially expressed genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue


A heatmap of all differentially expressed genes playing a potential regulatory role in the system (enriched in [TRANSPATH®](#) pathways) is presented in Figure 2.



Gene Expression Normalized by rows



Control: Non-tumour tissue
Experiment: Squamous Cell Carcinoma



- SET
- ANKK1
- MSL2
- STAG2
- CFP1
- EIF2AK1
- EXT2
- SMMT
- AFNKAP2
- ZNF1
- EXT1
- ODC2B8A
- LAMB2
- COLGALT1
- CD2H
- CHST1
- DLX3
- ITGA3
- ZNF164
- ANKK1
- SNAP70
- ALBA
- PRK8
- SLGA7
- TOP3
- TOP1B1
- TOP2B
- JARID1
- CEMP1
- CRTC3
- FBLN1
- PRPF11
- SPOPL
- PRF2
- ASF1A
- NAE1
- HI-2
- SNAIL2
- GPR1
- JARID1B
- ITGB4
- NECD
- HSP90A
- PRPF1
- ELP1
- QSOX1
- MN1
- SETD5
- BLDCL5B
- BEZ2
- TIMP1
- SNAPC1
- ZNF12
- SEPTIN7
- TULP4
- PHF2
- SNF2
- ANFY1
- TFEB
- NAAB1
- ILK
- COL1A4
- STK4
- PRK3
- CHST11
- FPSB4A
- CRIP1
- STAG1
- CFD
- MAF
- ARHGAP25
- SORL1
- LRRFRC
- EMF1
- EMIL4
- RFX
- NEK2
- LEB1D7
- PKP1
- RT1
- QSOX2
- HI-4
- CLIP1
- HACD1
- CAZ1
- PRK4
- LEP1
- PKC3
- PRK1
- PRK4
- DEK
- MPSNAP2
- ROCK1
- UBAP1
- TSG1
- YPS4B
- HLPDA
- EMC1
- FRS3B
- HMOX2
- GLI1
- ALDH3A1
- MCC1
- KLH8
- TRAF3
- SPRF3
- PRK1
- NECL1
- IFNA1
- ITSN2
- YPS4B
- EBF
- LEB1D1
- TNFSF2
- ELON4
- TOLLIP
- PELL1
- TNFR1
- BLVNB
- DLX1
- SLC1A1
- RASAL1
- LOC1
- ELON1
- FAP1
- SLK
- RASAL2
- FOXC2
- GJB2
- PRK1
- TNOC3
- DID2
- EPH1
- IL32
- IL18
- PLEC
- LAMA3
- USP4
- ITGB6
- IGFBP4
- COL1A1
- IGFBP7
- PRSS3
- COL6A2
- IGFBP5
- SPR4A
- IL13A1
- PRK1
- LOC1
- DKFZD4
- PRK1
- ILF1
- TNEM3
- SNX1
- NSD1
- SPR1B
- SYNE2
- ARL2
- PSME4
- PSME4
- GNS
- TPN4
- NF1
- NSD1A
- ACTB
- IGFBP2
- ITGB8
- TOP2A
- ILKAP1




Figure 2. Heatmap of genes enriched in Transpath categories. The colored bar at the top shows the types of the samples according to the legend in the upper right corner.

[See full diagram](#) →

Up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue:

768 significant up-regulated genes were taken for the mapping.

GO (biological process)



Figure 3. Enriched GO (biological process) of up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue.

[Full classification](#) →

TRANSPATH® Pathways (2024.1)




Figure 4. Enriched TRANSPATH® Pathways (2024.1) of up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue.

[Full classification →](#)

HumanPSD(TM) disease (2024.1)




Figure 5. Enriched HumanPSD(TM) disease (2024.1) of up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue. The size of the bars correspond to the number of biomarkers of the given disease found among the input set.

[Full classification](#) →

Down-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue:

650 significant down-regulated genes were taken for the mapping.

TRANSPATH® Pathways (2024.1)




Figure 7. Enriched TRANSPATH® Pathways (2024.1) of down-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue.

[Full classification →](#)

HumanPSD(TM) disease (2024.1)





Figure 8. Enriched HumanPSD(TM) disease (2024.1) of down-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue. The size of the bars correspond to the number of biomarkers of the given disease found among the input set.

[Full classification](#) →

The result of overall Gene Ontology (GO) analysis of the differentially expressed genes of the studied pathology can be summarized by the following diagram, revealing the most significant functional categories overrepresented among the observed (differentially expressed genes):



3.3. Analysis of enriched transcription factor binding sites and composite modules

In the next step a search for transcription factors binding sites (TFBS) was performed in the regulatory regions of the **target genes** by using the TF binding motif library of the [TRANSFAC®](#) database. We searched for so called **composite modules** that act as potential condition-specific **enhancers** of the **target genes** in their upstream regulatory regions (-1000 bp upstream of transcription start site (TSS)) and identify transcription factors regulating activity of the genes through such **enhancers**.

Classically, **enhancers** are defined as regions in the genome that increase transcription of one or several genes when inserted in either orientation at various distances upstream or downstream of the gene [8]. Enhancers typically have a length of several hundreds of nucleotides and are bound by multiple transcription factors in a cooperative manner [9].

In the current work, we use the Genomics data from the "Yes VCF track" track to predict positions of potential **enhancers** where the observed sequence variations may influence the gene expression in the pathology under study. We scan 5kb flanking regions and the body of all genes carrying the variations, with a sliding window of 1100bp size and find the position of the window with the maximal sum of the mutation weights, where we then perform the search for potential condition-specific enhancers (CMA model search).

We analyzed mutations that were revealed in the potential enhancers located upstream, downstream or inside the **target genes** (see Table 4). We identified 638 mutations potentially affecting gene regulation. Table 5 shows the following lists of PWMs whose sites were lost or gained due to these mutations. Weighting of mutations was done in respect to the significance of the change in TF affinity binding to the sequence. Mutations that maximally affected the change of binding affinity received higher weights. These PWMs were put in focus of the CMA algorithm that constructs the model of the enhancers by specifying combinations of TF motifs (see more details of the algorithm in the Methods section).

Table 4. Mutations revealed in Experiment: Squamous Cell Carcinoma versus Control: Non-tumour tissue

[See full table](#) →

ID	Gene symbol	Gene schematic representation	Number of variations
ENSG00000146648	EGFR		21
ENSG00000083857	FAT1		16
ENSG00000134871	COL4A2		13
ENSG00000186340	THBS2		10
ENSG00000226445	ENSG00000226445		9
ENSG00000145012	LPP		8
ENSG00000114999	TTL		7
ENSG00000142173	COL6A2		7
ENSG00000152291	TGOLN2		7
ENSG00000157214	STEAP2		7

Table 5. PWMs whose sites were lost or gained due to mutations in Experiment: Squamous Cell Carcinoma and Control: Non-tumour tissue

[See full table](#) →

ID	P-value (gains)	P-value (losses)	yesCount (gains)	yesCount (losses)
V\$EGR1_07	4.62E-2	1.48E-24	5	1134
V\$E2F7_04	3.9E-2	5.96E-23	11	744
V\$GLI2_05	2.5E-2	1.36E-22	11	2807
V\$E2F3_05	1.58E-2	3.85E-25	27	1467
V\$E2F1_Q4_01	1.5E-2	1.98E-27	11	1490
V\$TFCP2_06	2.67E-3	2.13E-16	7	3313
V\$RUNX3_01	5.83E-6	3.04E-24	151	1895
V\$E2F1_05	3.16E-7	6.77E-27	39	1042
V\$TEF_05	2.04E-7	1.43E-18	452	538
V\$MEIS1ELF1_01	2.29E-11	1.37E-16	2061	1805
V\$TFDP1_03	1.12E-12	6.17E-24	275	1398
V\$ZNF282_03	9.82E-20		803	
V\$MECP2_02	3.65E-20	1.39E-3	738	39
V\$SP1_09	3.12E-20	4.61E-2	342	4
V\$E2F1DP2_01	1.1E-20	1.22E-16	2155	2222
V\$E2F1EOMES_02	8.12E-21	5.89E-4	705	366
V\$SP2_09	1.61E-22	1.16E-2	236	25
V\$E2F3_10	7.25E-24	4.61E-2	1850	4
V\$RBAK_01	1.65E-24	5.74E-22	454	471
V\$E2F4_14	8.56E-25		1845	

We applied the Composite Module Analyst (CMA) [8] method to detect such potential enhancers, as targets of multiple TFs bound in a cooperative manner to the regulatory regions of the genes of interest. CMA applies a genetic algorithm to construct a generalized model of the enhancers by specifying combinations of TF motifs (from TRANSFAC®) whose sites are most frequently clustered together in the regulatory regions of the studied genes. CMA identifies the transcription factors that through their cooperation provide a synergistic effect and thus have a great influence on the gene regulation process.


Enhancer model potentially involved in regulation of target genes (up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue).

To build the most specific composite modules we choose top 300 significant up-regulated genes as the input of CMA algorithm. The obtained CMA model is then applied to compute CMA score for all up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue.


The model consists of 2 module(s). Below, for each module the following information is shown:

- PWMs producing matches,
- number of individual matches for each PWM,
- score of the best match.

Module 1:



Module 2:



Model score (-p*log10(pval)): 12.87

Wilcoxon p-value (pval): 3.95e-29

Penalty (p): 0.453

Average yes-set score: 2.98

Average no-set score: 2.00

AUC: 0.74

Separation point: 2.73

False-positive: 23.29%

False-negative: 38.80%




Table 6. List of top ten up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue with identified enhancers in their regulatory regions. **CMA score** - the score of the CMA model of the enhancer identified in the regulatory region.

[See full table](#) →

Ensembl IDs	Gene symbol	Gene description	CMA score	Factor names
ENSG00000089159	PXN	paxillin	6.88	ZNF281(h), CUX-1(h), TBX21(h), NF-E2L1(h), p53(h), p63(h), p73(h), SNAI1(h), SKIL(h), SMAD3(h)...
ENSG00000166165	CKB	creatine kinase B	6.53	SNAI1(h), CUX-1(h), TBX21(h), SKIL(h), ZNF281(h), NF-E2L1(h)
ENSG00000161203	AP2M1	adaptor related protein complex 2 subunit mu 1	6.34	p53(h), p63(h), p73(h), NF-E2L1(h), ZNF281(h), SKIL(h), JunD(h)
ENSG00000151150	ANK3	ankyrin 3	6.24	GATA-2(h), p53(h), p63(h), p73(h), SKIL(h), NF-E2L1(h), JunD(h), IRF-4(h)
ENSG00000196418	ZNF124	zinc finger protein 124	6.12	ZNF281(h), SNAI1(h), SMAD3(h), JunD(h), CUX-1(h), TBX21(h), GATA-2(h), p53(h), p63(h), p73(h)
ENSG00000122783	CYREN	cell cycle regulator of NHEJ	6.06	CUX-1(h), TBX21(h), JunD(h), NF-E2L1(h), p53(h), p63(h), p73(h), Hox-A10(h), PBX-4(h)
ENSG00000112234	FBXL4	F-box and leucine rich repeat protein 4	6.06	NF-E2L1(h), JunD(h), IRF-4(h), GATA-2(h), p53(h), p63(h), p73(h)
ENSG00000159335	PTMS	parathymosin	5.97	NF-E2L1(h), SKIL(h), SNAI1(h), JunD(h), GATA-2(h), p53(h), p63(h), p73(h)
ENSG00000111642	CHD4	chromodomain helicase DNA binding protein 4	5.86	ZNF281(h), SKIL(h), NF-E2L1(h), p53(h), p63(h), p73(h)
ENSG00000089693	MLF2	myeloid leukemia factor 2	5.79	SKIL(h), p53(h), p63(h), p73(h), IRF-4(h), ZNF281(h)


Enhancer model potentially involved in regulation of target genes (down-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue).

To build the most specific composite modules we choose top 300 significant down-regulated genes as the input of CMA algorithm. The obtained CMA model is then applied to compute CMA score for all down-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue.


The model consists of 2 module(s). Below, for each module the following information is shown:

- PWMs producing matches,
- number of individual matches for each PWM,
- score of the best match.

Module 1:



Module 2:



Model score (-p*log10(pval)): 16.58

Wilcoxon p-value (pval): 1.13e-35

Penalty (p): 0.475

Average yes-set score: 8.83

Average no-set score: 7.40

AUC: 0.76

Separation point: 7.74

False-positive: 36.75%

False-negative: 21.74%




Table 7. List of top ten down-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue with identified enhancers in their regulatory regions. **CMA score** - the score of the CMA model of the enhancer identified in the regulatory region.

[See full table](#) →

Ensembl IDs	Gene symbol	Gene description	CMA score	Factor names
ENSG00000188505	NCCRP1	NCCRP1, F-box associated domain containing	13.87	ARNT(h),ARNTL(h),ARNTL2(h),CLOCK(h), SMAD4(h), SMARCC2(h), ZNF282(h), TFII-I(h), GR(h), ZNF768(h)...
ENSG00000134954	ETS1	ETS proto-oncogene 1, transcription factor	13.28	ZNF768(h), SMARCC2(h), FOXA1(h), TFII-I(h), ARNT(h),ARNTL(h),ARNTL2(h),CLOCK(h), ZNF282(h), SMAD4(h)...
ENSG00000142327	RNPEPL1	arginyl aminopeptidase like 1	12.78	SMAD4(h), TFII-I(h), ZNF282(h), SMARCC2(h), ARNT(h),ARNTL(h),ARNTL2(h),CLOCK(h), FOXA1(h), ZNF768(h)...
ENSG00000078804	TP53INP2	tumor protein p53 inducible nuclear protein 2	12.76	SMARCC2(h), ZNF768(h), ARNT(h),ARNTL(h),ARNTL2(h),CLOCK(h), FOXA1(h), TFII-I(h), ZNF282(h), SMAD4(h)...
ENSG00000164442	CITED2	Cbp/p300 interacting transactivator with Glu/Asp rich carboxy-terminal domain 2	12.54	ARNT(h),ARNTL(h),ARNTL2(h),CLOCK(h), TFII-I(h), ZNF282(h), SMAD4(h), Sp1(h), FOXA1(h), ZNF768(h)
ENSG00000147676	MAL2	mal, T cell differentiation protein 2	12.52	SMARCC2(h), TFII-I(h), ZNF282(h), GR(h), SMAD4(h), Sp1(h), ZNF768(h)...
ENSG00000086548	CEACAM6	CEA cell adhesion molecule 6	12.38	FOXA1(h), ZNF768(h), ZNF282(h), SMAD4(h), TFII-I(h), GR(h), ARNT(h),ARNTL(h),ARNTL2(h),CLOCK(h)...
ENSG00000162368	CMPK1	cytidine/uridine monophosphate kinase 1	12.24	SMARCC2(h), TFII-I(h), SMAD4(h), ZNF282(h), ARNT(h),ARNTL(h),ARNTL2(h),CLOCK(h), FOXA1(h), ZNF768(h)
ENSG00000139289	PHLDA1	pleckstrin homology like domain family A member 1	12.12	ARNT(h),ARNTL(h),ARNTL2(h),CLOCK(h), TFII-I(h), SMAD4(h), ZNF282(h), Sp1(h), ZNF768(h), FOXA1(h)
ENSG00000182952	HMGN4	high mobility group nucleosomal binding domain 4	12.12	FOXA1(h), ZNF768(h), ZNF282(h), GR(h), SMAD4(h), TFII-I(h), ARNT(h),ARNTL(h),ARNTL2(h),CLOCK(h)

On the basis of the enhancer models we identified transcription factors potentially regulating the **target genes** of our interest. We found 17 and 13 transcription factors controlling expression of up- and down-regulated genes respectively (see Tables 8-9).

Table 8. Transcription factors of the predicted enhancer model potentially regulating the differentially expressed genes (up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue). **Yes-No ratio** is the ratio between frequencies of the sites in Yes sequences versus No sequences. It describes the level of the enrichment of binding sites for the indicated TF in the regulatory target regions. **Regulatory score** is the measure of involvement of the given TF in the controlling of expression of genes that encode master regulators presented below (through positive feedback loops).

[See full table](#) →


ID	Gene symbol	Gene description	Regulatory score	Yes-No ratio
MO000057832	SMAD3	SMAD family member 3	2.94	1.87
MO000042291	TP63	tumor protein p63	2.93	1.44
MO000019548	TP53	tumor protein p53	2.74	5.05
MO000044348	SNAI1	snail family transcriptional repressor 1	2.6	8.42
MO000024708	CUX1	cut like homeobox 1	2.47	3.03
MO000007834	JUND	JunD proto-oncogene, AP-1 transcription factor subunit	2.42	1.63
MO000089495	HOXA10	homeobox A10	2.38	5.61
MO000028475	SKIL	SKI like proto-oncogene	2.21	1.11
MO000028707	TP73	tumor protein p73	2.16	1.21
MO000028758	ZNF281	zinc finger protein 281	2.04	10.1

Table 9. Transcription factors of the predicted enhancer model potentially regulating the differentially expressed genes (down-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue). **Yes-No ratio** is the ratio between frequencies of the sites in Yes sequences versus No sequences. It describes the level of the enrichment of binding sites for the indicated TF in the regulatory target regions. **Regulatory score** is the measure of involvement of the given TF in the controlling of expression of genes that encode master regulators presented below (through positive feedback loops).

[See full table](#) →

ID	Gene symbol	Gene description	Regulatory score	Yes-No ratio
MO000019622	GTF2I	general transcription factor Iii	3.11	1.2
MO000033308	SP1	Sp1 transcription factor	2.85	1.93
MO000007834	JUND	JunD proto-oncogene, AP-1 transcription factor subunit	2.74	3.36
MO000031266	NR3C1	nuclear receptor subfamily 3 group C member 1	2.57	1.37
MO000020402	SMAD4	SMAD family member 4	2.56	1.77
MO000026696	ARNTL	aryl hydrocarbon receptor nuclear translocator like	2.33	1.29
MO000028681	CLOCK	clock circadian regulator	2.33	3.64
MO000114191	ARNT	aryl hydrocarbon receptor nuclear translocator	2.32	1.29
MO000177016	ZNF282	zinc finger protein 282	2.21	1.45
MO000026492	FOXA1	forkhead box A1	2.02	3.92

The following diagram represents the key transcription factors, which were predicted to be potentially regulating differentially expressed genes in the analyzed pathology: SMAD3, TP63, TP53, GTF2I, SP1 and JUND.



3.4. Finding master regulators in networks

In the second step of the upstream analysis common regulators of the revealed TFs were identified. We identified 10 signaling proteins whose structure and function is highly damaged by the mutations (see Table 10).

Table 10. Signaling proteins whose structure and function are damaged by the mutations in Experiment: Squamous Cell Carcinoma and Control: Non-tumour tissue

[See full table](#) →

ID	Title	Mutation count	Consequence	Codons
MO000208420	GJB3(h)	2	stop_gained	tGg/tAg
MO000109306	PSMA4(h)	1	stop_lost	Tga/Cga
MO000119197	wolframin(h)	1	stop_gained	Caa/Taa
MO000144222	APT2(h)	1	stop_lost	Tag/Cag
MO000172130	c3orf1(h)	1	NMD_transcript_variant,stop_lost	tGa/tCa
MO000175986	oas2(h)	1	stop_lost	tAg/tGg
MO000189841	ZSWIM1(h)	1	stop_gained	tGg/tAg
MO000212738	EMC10(h)	1	stop_lost	taG/taT
MO000219203	PSMG1(h)	1	NMD_transcript_variant,stop_lost	Taa/Caa
MO000222634	TCP11L1(h)	1	NMD_transcript_variant,stop_gained	Cag/Tag

Top 10 mutated proteins for Experiment: Squamous Cell Carcinoma and Control: Non-tumour tissue were used in the algorithm of master regulator search as a list of nodes of the signal transduction network that are removed from the network during the search of master regulators (see more details about the algorithm in the Methods section). These master regulators appear to be the key candidates for therapeutic targets as they have a master effect on regulation of intracellular pathways that activate the pathological process of our study. The identified master regulators are shown in Tables 11-12.

Table 11. Master regulators that may govern the regulation of **up-regulated** genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue. **Total rank** is the sum of the ranks of the master molecules sorted by keynode score, CMA score, transcriptomics data.

[See full table](#) →

ID	Master molecule name	Gene symbol	Gene description	logFC	Total rank
MO000016677	EGFR(h)	EGFR	epidermal growth factor receptor	4.44	112
MO000082228	EGFR-p60(h)	EGFR	epidermal growth factor receptor	4.44	123
MO000082230	EGFR-p110(h)	EGFR	epidermal growth factor receptor	4.44	123
MO000087397	EGFR-isoform4(h)	EGFR	epidermal growth factor receptor	4.44	123
MO000082277	EGFR-p170(h)	EGFR	epidermal growth factor receptor	4.44	126
MO000125420	EGFR(h){ub}n	EGFR	epidermal growth factor receptor	4.44	131
MO000042551	EGFR(h){pY}	EGFR	epidermal growth factor receptor	4.44	144
MO000199635	EGFR(h){pY869}{pY1016}{pY1092}	EGFR	epidermal growth factor receptor	4.44	155
MO000097590	EGFR(h){pY1016}	EGFR	epidermal growth factor receptor	4.44	156
MO000114808	EGFR(h){pY1197}	EGFR	epidermal growth factor receptor	4.44	156

Table 12. Master regulators that may govern the regulation of **down-regulated** genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue. **Total rank** is the sum of the ranks of the master molecules sorted by keynode score, CMA score, transcriptomics data.

[See full table](#) →

ID	Master molecule name	Gene symbol	Gene description	logFC	Total rank
MO000112152	MRG-1(h)	CITED2	Cbp/p300 interacting transactivator with Glu/Asp rich carboxy-terminal domain 2	-3.42	66
MO000056491	KAT2B(h)	KAT2B	lysine acetyltransferase 2B	-3.15	87
MO000033396	DUSP5(h)	DUSP5	dual specificity phosphatase 5	-4.8	154
MO000004672	ERK1(h)	MAPK3	mitogen-activated protein kinase 3	-2.26	173
MO000056883	ERK1-isoform1(h)	MAPK3	mitogen-activated protein kinase 3	-2.26	192
MO000256137	ERK1-isoform2(h)	MAPK3	mitogen-activated protein kinase 3	-2.26	200
MO000256138	ERK1-isoform3(h)	MAPK3	mitogen-activated protein kinase 3	-2.26	200
MO000137304	DUSP5(h)	DUSP5	dual specificity phosphatase 5	-4.8	208
MO000019174	Eck(h)	EPHA2	EPH receptor A2	-3.32	210
MO000112151	MRG-1-isoform1(h)	CITED2	Cbp/p300 interacting transactivator with Glu/Asp rich carboxy-terminal domain 2	-3.42	215

The intracellular regulatory pathways controlled by the above-mentioned master regulators are depicted in Figures 9 and 10. These diagrams display the connections between identified transcription factors, which play important roles in the regulation of differentially expressed genes, and selected master regulators, which are responsible for the regulation of these TFs.




Figure 9. Diagram of intracellular regulatory signal transduction pathways of up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue. Master regulators are indicated by red rectangles, transcription factors are blue rectangles, and green rectangles are intermediate molecules, which have been added to the network during the search for master regulators from selected TFs. Orange and blue frames highlight molecules that are encoded by up- and down-regulated genes, resp. [See full diagram](#) →




Figure 10. Diagram of intracellular regulatory signal transduction pathways of down-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue. Master regulators are indicated by red rectangles, transcription factors are blue rectangles, and green rectangles are intermediate molecules, which have been added to the network during the search for master regulators from selected TFs. Orange and blue frames highlight molecules that are encoded by up- and down-regulated genes, resp. [See full diagram](#) →

4. Finding prospective drug targets

The identified master regulators that may govern pathology associated genes were checked for druggability potential using HumanPSD™ [5] database of gene-disease-drug assignments and PASS [11-13] software for prediction of biological activities of chemical compounds on the basis of a (Q)SAR approach. Respectively, for each master regulator protein we have computed two Druggability scores: HumanPSD Druggability score and PASS Druggability score. Where Druggability score represents the number of drugs that are potentially suitable for inhibition (or activation) of the corresponding target either according to the information extracted from medical literature (from HumanPSD™ database) or according to cheminformatics predictions of compounds activity against the examined target (from PASS software).

The cheminformatics druggability check is done using a pre-computed database of spectra of biological activities of chemical compounds from a library of all small molecular drugs from HumanPSD™ database, 2507 pharmaceutically active known chemical compounds in total. The spectra of biological activities has been computed using the program PASS [11-13] on the basis of a (Q)SAR approach.

If both Druggability scores were below defined thresholds (see Methods section for the details) such master regulator proteins were not used in further analysis of drug prediction.

As a result we created the following two tables of prospective drug targets (top targets are shown here):



Table 13. Prospective drug targets selected from full list of identified master regulators filtered by Druggability score from *HumanPSD*TM database. **Druggability score** contains the number of drugs that are potentially suitable for inhibition (or activation) of the target. The drug targets are sorted according to the **Total rank** which is the sum of three ranks computed on the basis of the three scores: keynode score, CMA score and expression change score (logFC, if present). See Methods section for details.

[See full table](#) →

Gene symbol	Gene Description	Druggability score	logFC	Total rank
ITGA6	integrin subunit alpha 6	1	3.05	228
ODC1	ornithine decarboxylase 1	4	6.73	259
NTRK2	neurotrophic receptor tyrosine kinase 2	46	5.99	387
RBPJ	recombination signal binding protein for immunoglobulin kappa J region	1	1.81	619
TXNRD1	thioredoxin reductase 1	12	3.83	769
IGFBP5	insulin like growth factor binding protein 5	2	3.19	897



Table 14. Prospective drug targets selected from full list of identified master regulators filtered by Druggability score predicted by *PASS* software. Here, the **Druggability score** for master regulator proteins is computed as a sum of *PASS* calculated probabilities to be active as a target for various small molecular compounds. The drug targets are sorted according to the **Total rank** which is the sum of three ranks computed on the basis of the three scores: keynode score, CMA score and expression change score (logFC, if present). See Methods section for details.


[See full table](#) →

Gene symbol	Gene Description	Druggability score	logFC	Total rank
ITGA6	integrin subunit alpha 6	6.21	3.05	228
ITGB6	integrin subunit beta 6	6.21	3.05	228
ODC1	ornithine decarboxylase 1	1.05	6.73	259
NTRK2	neurotrophic receptor tyrosine kinase 2	12.4	5.99	387
PTPN12	protein tyrosine phosphatase non-receptor type 12	17.53	1.7	730
TXNRD1	thioredoxin reductase 1	4.89	3.83	769

Below we represent schematically the main mechanism of the studied pathology. In the schema we considered the top two drug targets of each of the two categories computed above. In addition we have added two top identified master regulators for which no drugs may be identified yet, but that are playing the crucial role in the molecular mechanism of the studied pathology. Thus the molecular mechanism of the studied pathology was predicted to be mainly based on the following key master regulators:

- integrins
- ornithine decarboxylase
- EGFR
- EGFR

This result allows us to suggest the following schema of affecting the molecular mechanism of the studied pathology:



Drugs which are shown on this schema: 1-(5-OXO-2,3,5,9B-TETRAHYDRO-1H-PYRROLO[2,1-A]ISOINDOL-9-YL)-3-(5-PYRROLIDIN-2-YL-1H-PYRAZOL-3-YL)-UREA, Terlipressin, Cytarabine, Erlotinib, Tegafur, Curcumin, Methyl aminolevulinic acid and Eflornithine, should be considered as a prospective research initiative for further drug repurposing and drug development. These drugs were selected as top matching treatments to the most prospective drug targets of the studied pathology, however, these results should be considered with special caution and are to be used for research purposes only, as there is not enough clinical information for adapting these results towards immediate treatment of patients.

The drugs given in dark red color on the schema are FDA approved drugs or drugs which have gone through various phases of clinical trials as active treatments against the selected targets.

The drugs given in pink color on the schema are drugs, which were cheminformatically predicted to be active against the selected targets.

5. Identification of potential drugs

In the last step of the analysis we strived to identify known activities as well as drugs with cheminformatically predicted activities that are potentially suitable for inhibition (or activation) of the identified molecular targets in the context of specified human diseases(s).

Proposed drugs are top ranked drug candidates, that were found to be active on the identified targets and were selected from 4 categories:

1. FDA approved drugs or used in clinical trials drugs for the studied pathology;
2. Repurposing drugs used in clinical trials for other pathologies;

3. Drugs, predicted by PASS to be active against identified drug targets and against the studied pathology;
4. Drugs, predicted by PASS to be active against identified drug targets but for other pathologies.

Proposed drugs were selected on the basis of Drug rank which was computed from the ranks sum based on the individual ranks of the following scores:

- Target activity score (depends on ranks of all targets that were found for the selected drug);
- Disease activity score (weighted sum of number of clinical trials on disease(s) under study where the selected drug is known to be applied or PASS Disease activity score - cheminformatically predicted property of the compound to be active against the studied disease(s));
- Clinical validity score (applicable only for drugs predicted on the basis of literature curation in HumanPSD™ database (Tables 16 and 17), reflects the number of the highest clinical trials phase on which the drug was tested for any pathology).

You can refer to the Methods section for more details on drug ranking procedure.

Based on the Drug rank, a numerical value of Drug score was calculated, which reflects the potential activity of the respective drug on the overall molecular mechanism of the studied pathology. Drug score values belong to the range from 1 to 100 and are calculated as a quotient of maximum drug rank and the drug rank of the given drug multiplied by 100.

Top drugs of each category are given in the tables below:

Drugs approved in clinical trials for Oncology



Table 15. Clinically approved (FDA, ENA, etc.) drugs for the studied pathology (most promising and clinically approved treatment candidates selected for the identified drug targets on the basis of literature curation in HumanPSD™ database)

[See full table](#) →

Name	Target names	Drug score	Disease activity score	Disease trial phase	Approved
Fluorouracil	PTPRC, BIRC5, CDKN1A	62	11	Phase 4: Carcinoma, Squamous Cell, Bowen's Disease, Breast Neoplasms, Carcinoma, Carcinoma, Basal Cell, Colorectal Neoplasms, Digestive System Neoplasms, Gastrointestinal Neoplasms, Glaucoma, Head and Neck Neoplasms, Hypopigmentation, Intestinal Neoplasms, Keratosis, Keratosis, Actinic, Neoplasms, Neoplasms, Basal Cell, Neoplasms, Squamous Cell, Photosensitivity Disorders, Postoperative Complications, Skin Diseases, Skin Neoplasms, Squamous Cell Carcinoma of Head and Neck, Vitiligo	Carcinoma, Squamous Cell (ClinicalTrials, ClinicalTrials, ClinicalTrials)

The **Disease trial phase** column reflects the maximum clinical trials phase in which the drug was studied for the analyzed pathology.

Drugs approved in clinical trials



Table 16. Drugs used in clinical trials for the studied pathology (most promising treatment candidates selected for the identified drug targets on the basis of literature curation in [HumanPSD™](#) database)

[See full table](#) →

Name	Target names	Drug score	Disease activity score	Disease trial phase
Erlotinib	RPS6KA3, MAP4K4, MET, NEK7, PAK2, GSK3B, LATS1, MAP3K11, PRKAA1, BMPR2, ILK, EGFR, SYK, EIF2AK2, BIRC5, IGF1R, JAK1, TGFB1, SRC, CSNK2A2, PRKD3, PIP4K2B, NTRK2, CSNK1G2, EPHB4, YES1, BRAF, STK11, STK4, PTK2, PIK3CA, LYN, CSNK2A1, PRKACB, CSF1R	99	7	Phase 3: Carcinoma, Squamous Cell, Brain Neoplasms, Carcinoma, Carcinoma, Adenosquamous, Carcinoma, Hepatocellular, Carcinoma, Large Cell, Carcinoma, Non-Small-Cell Lung, Head and Neck Neoplasms, Lung Neoplasms, Neoplasm Metastasis, Neoplasms, Thoracic Neoplasms
Tegafur	ITGA6, VEGFA, ITGB5, EGFR, ITGB1, PTK2, ITGB4, ITGA3	99	5	Phase 3: Carcinoma, Squamous Cell, Adenocarcinoma, Ascites, Carcinoma, Cholangiocarcinoma, Esophageal Neoplasms, Esophageal Squamous Cell Carcinoma, Nasopharyngeal Carcinoma, Neoplasm Metastasis, Neoplasms, Pancreatic Neoplasms, Stomach Neoplasms
Gefitinib	RPS6KA3, MAP4K4, MET, NEK7, PAK2, GSK3B, LATS1, MAP3K11, PRKAA1, BMPR2, EGFR, SYK, EIF2AK2, IGF1R, JAK1, TGFB1, SRC, CSNK2A2, PRKD3, PIP4K2B, NTRK2, CSNK1G2, EPHB4, YES1, BRAF, STK11, STK4, PTK2, PIK3CA, LYN, CSNK2A1, PRKACB, CSF1R	95	3	Phase 2: Carcinoma, Squamous Cell, Adenocarcinoma, Adenocarcinoma of Lung, Brain Neoplasms, Breast Neoplasms, Carcinoid Tumor, Carcinoma, Carcinoma, Islet Cell, Carcinoma, Non-Small-Cell Lung, Carcinoma, Renal Cell, Carcinoma, Transitional Cell, Fallopian Tube Neoplasms, Gastrinoma, Gastrointestinal Neoplasms, Glioblastoma, Glucagonoma, Head and Neck Neoplasms, Insulinoma, Intestinal Neoplasms, Lung Diseases, Lung Neoplasms, Malignant Carcinoid Syndrome, Mesothelioma, Mesothelioma, Malignant, Nasopharyngeal Carcinoma, Nasopharyngeal Neoplasms, Neoplasm Metastasis, Neoplasms, Neoplasms, Squamous Cell, Neuroblastoma, Neuroendocrine Tumors, Ovarian Neoplasms, Peritoneal Neoplasms, Recurrence, Respiratory Tract Diseases, Respiratory Tract Neoplasms, Sarcoma, Sarcoma, Synovial, Somatostatinoma, Squamous Cell Carcinoma of Head and Neck, Stomach Neoplasms, Thoracic Neoplasms, Urinary Bladder Neoplasms, Vipoma
Lapatinib	RPS6KA3, MAP4K4, MET, NEK7, PAK2, GSK3B, LATS1, MAP3K11, PRKAA1, BMPR2, EGFR, SYK, EIF2AK2, BIRC5, IGF1R, JAK1, TGFB1, SRC, CSNK2A2, PRKD3, PIP4K2B, NTRK2, CSNK1G2, EPHB4, YES1, BRAF, STK11, STK4, PTK2, PIK3CA, LYN, CSNK2A1, PRKACB, CSF1R	94	2	Phase 2: Carcinoma, Squamous Cell, Adenocarcinoma, Brain Neoplasms, Breast Diseases, Breast Neoplasms, Carcinoma, Carcinoma, Ductal, Carcinoma, Ductal, Breast, Carcinoma, Non-Small-Cell Lung, Carcinoma, Small Cell, Carcinoma, Transitional Cell, Fibroma, Glioblastoma, Glioma, Gliosarcoma, Head and Neck Neoplasms, Liver Neoplasms, Neoplasm Metastasis, Neoplasms, Neurilemmoma, Neuroblastoma, Neurofibroma, Neurofibromatoses, Neurofibromatosis 1, Neurofibromatosis 2, Neuroma, Neuroma, Acoustic, Sarcoma, Small Cell Lung Carcinoma, Squamous Cell Carcinoma of Head and Neck, Thymoma, Urinary Bladder Neoplasms

Vandetanib	RPS6KA3, MAP4K4, MET, NEK7, PAK2, GSK3B, LATS1, MAP3K11, PRKAA1, BMPR2, EGFR, SYK, EIF2AK2, IGF1R, JAK1, VEGFA, TGFBR1, SRC, CSNK2A2, PRKD3, PIP4K2B, NTRK2, CSNK1G2, EPHB4, YES1, BRAF, STK11, STK4, PTK2, PIK3CA, LYN, CSNK2A1, PRKACB, CSF1R	94	2	Phase 2: Carcinoma, Squamous Cell, Astrocytoma, Brain Abscess, Breast Neoplasms, Carcinoma, Carcinoma, Hepatocellular, Carcinoma, Neuroendocrine, Carcinoma, Non-Small-Cell Lung, Carcinoma, Transitional Cell, Endocrine Gland Neoplasms, Fallopian Tube Neoplasms, Glioblastoma, Glioma, Gliosarcoma, Lung Neoplasms, Mesothelioma, Mesothelioma, Malignant, Multiple Endocrine Neoplasia, Multiple Endocrine Neoplasia Type 2a, Multiple Endocrine Neoplasia Type 2b, Multiple Myeloma, Neoplasm Metastasis, Neoplasms, Neoplasms, Plasma Cell, Oligodendroglioma, Ovarian Neoplasms, Peritoneal Neoplasms, Pleural Effusion, Pleural Effusion, Malignant, Sarcoma, Squamous Cell Carcinoma of Head and Neck, Thyroid Diseases, Thyroid Neoplasms, Urinary Bladder Neoplasms
------------	---	----	---	---

The ***Disease trial phase*** column reflects the maximum clinical trials phase in which the drug was studied for the analyzed pathology.

Repurposing drugs



Table 17. Repurposed drugs used in clinical trials for other pathologies (prospective drugs against the identified drug targets on the basis of literature curation in [HumanPSD™](#) database)

[See full table](#) →

Name	Target names	Drug score	Maximum trial phase
Curcumin	CD44, CDK6, MET, SMAD2, SMAD3, HSPA5, HK2, CCNB1, GSK3B, VIM, CDK4, PTEN, PRKAA1, CDC20, PARP1, EGFR, CTNNB1, SLC2A1, HIF1A, ATM, MAP1LC3B, BIRC5, SUZ12, JAK1, ATR, VEGFA, CEBPA, PCNA, YWHAE, EIF2S1, APP, NEDD4, TFRC, EPAS1, PSEN1, BIRC2, MMP14, CCND1, SKP2, IGFBP5, SMO, CCNA2, HMGB1, CDKN1A, JAG1	91	Phase 4: Cardiovascular Abnormalities, Cysts, Diabetes Mellitus, Diabetes Mellitus, Type 2, Glucose Intolerance, Insulin Resistance, Irritable Bowel Syndrome, Kidney Diseases, Kidney Diseases, Cystic, Periodontitis, Polycystic Kidney Diseases, Polycystic Kidney, Autosomal Dominant, Prediabetic State, Syndrome
Flavopiridol	RPS6KA3, CDK6, MAP4K4, MET, NEK7, PAK2, GSK3B, LATS1, MAP3K11, CDK4, PRKAA1, BMPR2, EGFR, SYK, EIF2AK2, IGF1R, JAK1, TGFBR1, SRC, CSNK2A2, PRKD3, PIP4K2B, NTRK2, CSNK1G2, EPHB4, YES1, BRAF, STK11, PIK3CB, STK4, PTK2, PIK3CA, LYN, CSNK2A1, PRKACB, CSF1R	91	Phase 2: Embolism, Head and Neck Neoplasms, Lymphoma, Lymphoma, B-Cell, Lymphoma, Large B-Cell, Diffuse, Lymphoma, Mantle-Cell, Lymphoma, Non-Hodgkin, Neoplasms, Sarcoma, Thromboembolism
seliciclib	RPS6KA3, ROCK2, MAP4K4, MET, NEK7, PAK2, GSK3B, LATS1, MAP3K11, CDK4, PRKAA1, BMPR2, EGFR, SYK, SGK1, EIF2AK2, IGF1R, JAK1, TGFBR1, SRC, CSNK2A2, PRKD3, PIP4K2B, NTRK2, CSNK1G2, EPHB4, YES1, BRAF, STK11, STK4, PTK2, PIK3CA, LYN, CSNK2A1, PRKACB, CSF1R	91	Phase 2: Cystic Fibrosis, Cysts, Fibrosis
Tofacitinib	RPS6KA3, ROCK2, MAP4K4, MET, NEK7, PAK2, GSK3B, LATS1, MAP3K11, PRKAA1, BMPR2, EGFR, SYK, SGK1, EIF2AK2, IGF1R, JAK1, TGFBR1, SRC, CSNK2A2, PRKD3, PIP4K2B, NTRK2, CSNK1G2, EPHB4, YES1, BRAF, STK11, STK4, PTK2, PIK3CA, LYN, CSNK2A1, PRKACB, CSF1R	91	Phase 4: Alopecia, Alopecia Areata, Aortic Arch Syndromes, Arteritis, Arthritis, Arthritis, Psoriatic, Arthritis, Rheumatoid, Axial Spondyloarthritis, COVID-19, Colitis, Colitis, Ulcerative, Crohn Disease, Embolism, Granuloma, Granulomatosis with Polyangiitis, Inflammatory Bowel Diseases, Intestinal Diseases, Necrosis, ST Elevation Myocardial Infarction, Spondylarthritis, Spondylitis, Systemic Vasculitis, Takayasu Arteritis, Thromboembolism, Ulcer, Vasculitis
1-(5-Tert-Butyl-2-P-Tolyl-2h-Pyrazol-3-Yl)-3-[4-(2-Morpholin-4-Yl-Ethoxy)-Naphthalen-1-Yl]-Urea	RPS6KA3, ROCK2, MAP4K4, MET, NEK7, PAK2, GSK3B, LATS1, MAP3K11, PRKAA1, BMPR2, EGFR, SYK, SGK1, EIF2AK2, IGF1R, JAK1, TGFBR1, SRC, CSNK2A2, PRKD3, PIP4K2B, NTRK2, CSNK1G2, EPHB4, YES1, BRAF, STK11, STK4, PTK2, PIK3CA, LYN, CSNK2A1, PRKACB, CSF1R	91	Phase 2: Arthritis, Arthritis, Rheumatoid, Psoriasis

The **Maximum trial phase** column reflects the maximum clinical trials phase in which the drug was studied for any pathology.



No prospective drugs were found, which would be predicted by PASS software to be active against the identified drug targets and would be predicted to have biological activity against the studied disease(s).



Table 18. Prospective drugs, predicted by PASS software to be active against the identified drug targets, though without cheminformatically predicted activity against the studied disease(s) (drug candidates predicted with the cheminformatics tool PASS)

[See full table](#) →

Name	Target names	Drug score	Target activity score
{(2Z)-4-AMINO-2-[(4-METHOXYPHENYL)IMINO]-2,3-DIHYDRO-1,3-THIAZOL-5-YL}(4-METHOXYPHENYL)METHANONE	CCND1, CDK6, CCND3, CCNB1, CCNA2, CCNT1, CCNB2, CDK4	100	4.63
3-Bromo-7-Nitroindazole	RPS6KA3, CCND1, CDK6, HSPD1, CCND3, CCNB1, GSK3B, CCNA2, CCNT1, CCNB2, CDK4	100	3.29
O6-CYCLOHEXYLMETHOXY-2-(4'-SULPHAMOYLANILINO) PURINE	CCND1, CDK6, CCND3, CCNB1, CCNA2, CCNT1, CDK4, CCNB2	100	3.03
2-ANILINO-6-CYCLOHEXYLMETHOXPURINE	CCND1, CDK6, CCND3, CCNB1, CCNA2, CCNT1, CDK4, CCNB2	100	2.85
2-(2-HYDROXYETHYLAMINO)-6-(3-CHLOROANILINO)-9-ISOPROPYLPURINE	CCND1, CDK6, SRC, CCND3, CCNB1, CCNA2, CCNT1, CDK4, CCNB2	100	2.79

As the result of drug search we propose the following drugs as most promising candidates for treating the pathology under study: Erlotinib, Curcumin and {(2Z)-4-AMINO-2-[(4-METHOXYPHENYL)IMINO]-2,3-DIHYDRO-1,3-THIAZOL-5-YL}(4-METHOXYPHENYL)METHANONE. These drugs were selected for acting on the following targets: NTRK2, IGFBP5 and CCND1, which were predicted to be active in the molecular mechanism of the studied pathology.

The selected drugs are top ranked drug candidates from each of the four categories of drugs: (1) FDA approved drugs or used in clinical trials drugs for the studied pathology; (2) repurposing drugs used in clinical trials for other pathologies; (3) drugs, predicted by PASS software to be active against the studied pathology; (4) drugs, predicted by PASS software to be repurposed from other pathologies.

Supplementary drug info

In addition to the approved and repurposed drugs proposed by Genome Enhancer, below the **Supplementary drug info** table is given, which contains an extended list of drugs used for treatment of neoplasms. Those drugs which were predicted by Genome Enhancer as prospective treatment candidates for the studied case (both approved and repurposed) have a respective **Predicted Drug Score** assigned to them. This value on a scale from 1 to 100 reflects the potential activity of the respective drug on the overall molecular mechanism of the studied pathology. The **Predicted Drug Score** column contains the "-" (Not Identified) value in case the drug targets of the respective treatment were not found in the molecular mechanism of the studied pathology.

Table 19. Supplementary drug info: extended list of drugs used for treatment of neoplasms with respective drug scores predicted for the studied pathology.

Drug	Disease	Predicted Drug Score
Abarelix	Prostatic Neoplasms	-
Abemaciclib	Breast Neoplasms	86
Abiraterone	Prostatic Neoplasms, Castration-Resistant	-
Abiraterone acetate	Prostatic Neoplasms, Castration-Resistant	-

Acalabrutinib	Lymphoma, Mantle-Cell	-
Acitretin	Psoriasis	-
Ado-trastuzumab emtansine	Breast Neoplasms Neoplasms	36
Afatinib	Carcinoma, Non-Small-Cell Lung	56
Aflibercept	Colorectal Neoplasms Diabetic Retinopathy Edema Vascular Diseases Wet Macular Degeneration	20
Alectinib	Carcinoma, Non-Small-Cell Lung	10
Alemtuzumab	Brain Abscess Leukemia, Lymphocytic, Chronic, B-Cell Multiple Sclerosis Multiple Sclerosis, Relapsing-Remitting Sclerosis	-
Alitretinoin	Sarcoma, Kaposi	-
Alpelisib	Breast Neoplasms	82
Altretamine	Ovarian Neoplasms	-
Aminolevulinic acid	Keratosis Keratosis, Actinic	-
Anagrelide	Thrombocythemia, Essential Thrombocytosis	-
Anastrozole	Breast Neoplasms Hypersensitivity Obesity Obesity, Morbid Recurrence Weight Loss	-
Apalutamide	Prostatic Neoplasms, Castration-Resistant	-
Aprepitant	Nausea Neoplasms Postoperative Nausea and Vomiting	-
Arsenic trioxide	Leukemia, Promyelocytic, Acute	85
Atezolizumab	Carcinoma, Non-Small-Cell Lung Carcinoma, Transitional Cell Triple Negative Breast Neoplasms	-
Avelumab	Carcinoma, Merkel Cell Carcinoma, Renal Cell Carcinoma, Transitional Cell	-
Axitinib	Carcinoma, Renal Cell	-
Azacitidine	Anemia, Refractory Anemia, Refractory, with Excess of Blasts Leukemia, Myelomonocytic, Chronic Myelodysplastic Syndromes Preleukemia Syndrome	2
Belinostat	Lymphoma, T-Cell, Peripheral	36
Bendamustine	Leukemia, Lymphocytic, Chronic, B-Cell Leukemia, Lymphoid	-
Bevacizumab	Breast Neoplasms Colonic Neoplasms Colorectal Neoplasms Corneal Neovascularization Diabetic Retinopathy Dilatation, Pathologic Edema Epistaxis Glaucoma Hemorrhage Macular Degeneration Macular Edema Neoplasm Metastasis Neoplasms Neovascularization, Pathologic Optic Nerve Diseases Pterygium Rectal Neoplasms Retinal Detachment Retinal Diseases Retinal Vein Occlusion Telangiectasia, Hereditary Hemorrhagic Telangiectasis Vitreous Hemorrhage	14
Bexarotene	Lymphoma, T-Cell Lymphoma, T-Cell, Cutaneous	-
Bicalutamide	Prostatic Neoplasms	7
Binimetinib	Melanoma	-
Blinatumomab	Precursor B-Cell Lymphoblastic Leukemia-Lymphoma	-
Bortezomib	Brain Abscess Glomerulonephritis Glomerulonephritis, IGA Kidney Diseases Multiple Myeloma Neoplasms, Plasma Cell Nephritis Renal Insufficiency	37
Bosutinib	Leukemia, Myelogenous, Chronic, BCR-ABL Positive	61
Brentuximab vedotin	Hodgkin Disease Lymphoma Lymphoma, Large-Cell, Anaplastic Lymphoma, T-Cell, Peripheral	-
Brigatinib	Carcinoma, Non-Small-Cell Lung	70
Buserelin	Prostatic Neoplasms	-
Cabazitaxel	Prostatic Neoplasms, Castration-Resistant	12
Cabergoline	Drug-Related Side Effects and Adverse Reactions Pituitary Neoplasms	-
Cabozantinib	Thyroid Neoplasms	78
Capecitabine	Breast Neoplasms Colonic Neoplasms Colorectal Neoplasms	-
Carboplatin	Carcinoma, Non-Small-Cell Lung Lung Neoplasms Neoplasms Neuroendocrine Tumors Ovarian Neoplasms Retinoblastoma	-
Carfilzomib	Multiple Myeloma	8
Carmustine	Astrocytoma Glioblastoma Hodgkin Disease Medulloblastoma Multiple Myeloma Neoplasms	-

Ceritinib	Carcinoma, Non-Small-Cell Lung	55
Cetuximab	Colorectal Neoplasms	43
Cinacalcet	Anemia Calcinosis Cardiovascular Diseases Hyperparathyroidism Hyperparathyroidism, Secondary Kidney Diseases Kidney Failure, Chronic Neoplasm Metastasis Neoplasms Parathyroid Neoplasms Renal Insufficiency Vascular Calcification Vascular Diseases Vision Disorders	-
Cisplatin	Carcinoma, Squamous Cell Neoplasms Uterine Cervical Neoplasms Carcinoma, Non-Small-Cell Lung Esophageal Neoplasms Carcinoma	-
Cladribine	Leukemia, Hairy Cell	34
Clofarabine	Precursor Cell Lymphoblastic Leukemia-Lymphoma	59
Cobimetinib	Melanoma	-
Copanlisib	Lymphoma, Follicular	71
Crizotinib	Carcinoma, Non-Small-Cell Lung	93
Cyproterone acetate	Prostatic Neoplasms	-
Dabrafenib	Melanoma	9
Dacomitinib	Carcinoma, Non-Small-Cell Lung	72
Daratumumab	Multiple Myeloma	-
Dasatinib	Leukemia, Myelogenous, Chronic, BCR-ABL Positive Leukemia, Myeloid, Chronic-Phase Precursor Cell Lymphoblastic Leukemia-Lymphoma	88
Decitabine	Anemia, Refractory Anemia, Refractory, with Excess of Blasts Leukemia, Myelomonocytic, Chronic Myelodysplastic Syndromes	-
Degarelix	Cardiovascular Diseases Prostatic Neoplasms Vascular Diseases	-
Denosumab	Arthritis, Rheumatoid Bone Diseases Bone Diseases, Metabolic Breast Neoplasms Hyperparathyroidism Hyperparathyroidism, Primary Metabolic Diseases Neoplasm Metastasis Neoplasms Osteoporosis Osteoporosis, Postmenopausal Prostatic Neoplasms	-
Dexrazoxane	Breast Neoplasms Cardiomyopathies	70
Dienogest	Menorrhagia	-
Dinutuximab	Neuroblastoma	-
Docetaxel	Breast Neoplasms Carcinoma, Non-Small-Cell Lung Prostatic Neoplasms Squamous Cell Carcinoma of Head and Neck Stomach Neoplasms	-
Doxorubicin	Neoplasms Multiple Myeloma Carcinoma, Ovarian Epithelial Ovarian Neoplasms Leukemia, Lymphoid Breast Neoplasms Lymphoma, Follicular Thyroid Neoplasms Triple Negative Breast Neoplasms Glioma	87
Durvalumab	Carcinoma, Non-Small-Cell Lung Carcinoma, Transitional Cell	-
Dutasteride	Alcoholism Hyperplasia Hypertrophy Neoplasms Prostatic Hyperplasia	-
Duvelisib	Leukemia, Lymphocytic, Chronic, B-Cell Lymphoma, Follicular	-
Elotuzumab	Multiple Myeloma	-
Enasidenib	Leukemia, Myeloid, Acute	-
Encorafenib	Colorectal Neoplasms Melanoma	3
Enfortumab vedotin	Carcinoma, Transitional Cell Neoplasms	-
Entrectinib	Carcinoma, Non-Small-Cell Lung	63
Enzalutamide	Prostatic Neoplasms Prostatic Neoplasms, Castration-Resistant	-
Epirubicin	Breast Neoplasms	78
Erdafitinib	Urinary Bladder Neoplasms	65
Eribulin	Breast Neoplasms Drug-Related Side Effects and Adverse Reactions Neoplasms	-
Erlotinib	Carcinoma, Non-Small-Cell Lung Neoplasms Pancreatic Neoplasms	99
Erlotinib hydrochloride	Carcinoma, Non-Small-Cell Lung Gastrointestinal Stromal Tumors	-
Estramustine	Prostatic Neoplasms	-
Ethinyl Estradiol	Acne Vulgaris Neoplasms	13
Everolimus	Angiomyolipoma Arthrogyrosis Astrocytoma Breast Neoplasms Carcinoma, Renal Cell Cysts Idiopathic Pulmonary Fibrosis Kidney Diseases, Cystic Kidney Failure, Chronic Lipoma Neuroendocrine Tumors Primary Graft Dysfunction Sclerosis Tuberous Sclerosis	72
Exemestane	Breast Neoplasms	-

Fedratinib	Primary Myelofibrosis	-
Finasteride	Hyperplasia Neoplasms Prostatic Hyperplasia	-
Flavopiridol	Leukemia, Lymphocytic, Chronic, B-Cell	91
Fluorouracil	Skin Neoplasms Neoplasms, Basal Cell Neoplasms, Second Primary Neoplasms, Squamous Cell Neoplasms Colorectal Neoplasms Pancreatic Neoplasms	62
Flooxymesterone	Breast Neoplasms Hypogonadism Puberty, Delayed	22
Flutamide	Premenstrual Dysphoric Disorder Premenstrual Syndrome Prostatic Neoplasms	56
Fulvestrant	Breast Neoplasms	-
Gefitinib	Carcinoma, Non-Small-Cell Lung	95
Gemcitabine	Breast Neoplasms Carcinoma, Non-Small-Cell Lung Ovarian Neoplasms Pancreatic Neoplasms	35
Gemtuzumab ozogamicin	Leukemia, Myeloid, Acute	-
Gilteritinib	Leukemia, Myeloid, Acute	55
Glasdegib	Leukemia, Myeloid, Acute	67
Goserelin	Atrophy Breast Neoplasms Bulbo-Spinal Atrophy, X-Linked Endometriosis Muscular Atrophy Myoma Prostatic Neoplasms	-
Histrelin	Puberty, Precocious	-
Homoharringtonine	Leukemia, Myelogenous, Chronic, BCR-ABL Positive	58
Ibritumomab	Lymphoma, B-Cell Lymphoma, Follicular	-
Ibrutinib	Graft vs Host Disease Leukemia, Lymphocytic, Chronic, B-Cell Lymphoma, B-Cell, Marginal Zone Lymphoma, Mantle-Cell Waldenstrom Macroglobulinemia	14
Idarubicin	Leukemia, Myeloid, Acute	44
Idelalisib	Leukemia, Lymphocytic, Chronic, B-Cell Lymphoma, Follicular	11
Ifosfamide	Neoplasms	-
Imatinib	Leukemia, Myelogenous, Chronic, BCR-ABL Positive Mastocytosis, Systemic Neoplasms	91
Inotuzumab ozogamicin	Precursor B-Cell Lymphoblastic Leukemia-Lymphoma	-
Ipilimumab	Carcinoma, Renal Cell Melanoma	-
Irinotecan	Colorectal Neoplasms	70
Ivosidenib	Leukemia, Myeloid, Acute	-
Ixabepilone	Breast Neoplasms	-
Ixazomib	Multiple Myeloma	-
Lapatinib	Breast Neoplasms	94
Larotrectinib	Neoplasm Metastasis	75
Lenalidomide	Brain Abscess Lupus Erythematosus, Cutaneous Myelodysplastic Syndromes Neoplasms, Plasma Cell	12
Lenvatinib	Carcinoma, Hepatocellular Carcinoma, Renal Cell Thyroid Neoplasms	-
Letrozole	Breast Neoplasms Cysts Fibroma Myofibroma Myoma Ovarian Cysts Syndrome	-
Leuprolide	Hot Flashes Ovarian Hyperstimulation Syndrome Prostatic Neoplasms Puberty, Precocious	-
Levamisole	Ascariasis Colonic Neoplasms Helminthiasis	-
Levonorgestrel	Epilepsy Hyperplasia Menorrhagia	-
Lomustine	Brain Neoplasms Hodgkin Disease	-
Lonafarnib	Breast Neoplasms Carcinoma, Non-Small-Cell Lung Central Nervous System Neoplasms Colorectal Neoplasms Gliosarcoma Head and Neck Neoplasms Leukemia, Myelomonocytic, Chronic Liver Neoplasms Lymphoma Myelodysplastic Syndromes Ovarian Neoplasms Urethral Neoplasms Urinary Bladder Neoplasms	51
Lorlatinib	Carcinoma, Non-Small-Cell Lung	83
Masoprocol	Keratosis, Actinic	-
Medroxyprogesterone Acetate	Depression Depression, Postpartum Depressive Disorder Metrorrhagia Neoplasms Uterine Hemorrhage	12
Megestrol acetate	Acquired Immunodeficiency Syndrome Bites and Stings Breast Neoplasms Pain Wasting Syndrome	15
Methotrexate	Neoplasms Breast Neoplasms Head and Neck Neoplasms Ovarian Neoplasms Lymphoma, T-Cell, Peripheral Brain Neoplasms Colorectal	50

	Neoplasms Neuroblastoma Carcinoma, Squamous Cell	
Methyltestosterone	Breast Neoplasms Hypogonadism Puberty, Delayed	-
Midostaurin	Leukemia, Mast-Cell Leukemia, Myeloid, Acute Mastocytosis, Systemic	90
Mitotane	Adrenocortical Carcinoma	-
Mitoxantrone	Autoimmune Diseases Autoimmune Diseases of the Nervous System Demyelinating Autoimmune Diseases, CNS Immune System Diseases Leukemia, Myeloid, Acute Multiple Sclerosis Myelitis Myelitis, Transverse Nervous System Diseases Neuromyelitis Optica Prostatic Neoplasms, Castration-Resistant	53
Mogamulizumab	Mycosis Fungoides Neoplasms Sezary Syndrome	-
Moxetumomab pasudotox	Leukemia, Hairy Cell Neoplasms	-
Necitumumab	Carcinoma, Non-Small-Cell Lung Neoplasms	-
Nelarabine	Precursor T-Cell Lymphoblastic Leukemia-Lymphoma	-
Neratinib	Breast Neoplasms	67
Nilotinib	Blast Crisis Leukemia, Myelogenous, Chronic, BCR-ABL Positive Leukemia, Myeloid, Chronic-Phase	17
Nilutamide	Prostatic Neoplasms	-
Nintedanib	Fibrosis Idiopathic Pulmonary Fibrosis	51
Niraparib	Carcinoma, Ovarian Epithelial Fallopian Tube Neoplasms Peritoneal Neoplasms	54
Nivolumab	Carcinoma, Non-Small-Cell Lung Kidney Neoplasms Neoplasms Lung Neoplasms Melanoma	-
Obinutuzumab	Leukemia, Lymphocytic, Chronic, B-Cell	-
Octreotide	Acromegaly Adenoma Ascites Carcinoid Tumor Fistula Pancreatic Fistula Pituitary Diseases Renal Insufficiency Vipoma	4
Ofatumumab	Leukemia, Lymphocytic, Chronic, B-Cell	-
Olaparib	Breast Neoplasms Carcinoma, Ovarian Epithelial Fallopian Tube Neoplasms Ovarian Neoplasms Pancreatic Neoplasms Peritoneal Neoplasms Prostatic Neoplasms, Castration-Resistant	72
Olaratumab	Sarcoma	-
Osimertinib	Carcinoma, Non-Small-Cell Lung	24
Oxaliplatin	Colonic Neoplasms Colorectal Neoplasms Neoplasms Rectal Neoplasms	59
Paclitaxel	Acute Coronary Syndrome Angina Pectoris Arteriosclerosis Breast Neoplasms Carcinoma, Non-Small-Cell Lung Cardiovascular Diseases Coronary Artery Disease Coronary Disease Coronary Stenosis Heart Diseases Myocardial Ischemia Ovarian Neoplasms Vascular Diseases	66
Palbociclib	Breast Neoplasms	69
Panitumumab	Colorectal Neoplasms	81
Panobinostat	Multiple Myeloma	-
Pazopanib	Carcinoma Carcinoma, Renal Cell Sarcoma	89
Pembrolizumab	Carcinoma, Hepatocellular Carcinoma, Merkel Cell Carcinoma, Non-Small-Cell Lung Carcinoma, Renal Cell Carcinoma, Transitional Cell Hodgkin Disease Melanoma Neoplasms Stomach Neoplasms	-
Pemetrexed	Carcinoma, Non-Small-Cell Lung Mesothelioma	-
Pentostatin	Leukemia, Hairy Cell	38
Pertuzumab	Breast Neoplasms	-
Pomalidomide	Multiple Myeloma	14
Ponatinib	Leukemia, Myelogenous, Chronic, BCR-ABL Positive Precursor Cell Lymphoblastic Leukemia-Lymphoma	82
Pralatrexate	Lymphoma, T-Cell, Peripheral	-
Radium Ra 223 Dichloride	Prostatic Neoplasms, Castration-Resistant	-
Ramucirumab	Stomach Neoplasms	-
Rasburicase	Hyperuricemia Leukemia Lymphoma Neoplasms Syndrome Tumor Lysis Syndrome	-
Regorafenib	Colorectal Neoplasms	29
Relugolix	Prostatic Neoplasms	-
Ribociclib	Breast Neoplasms	77

Rituximab	Arthritis Arthritis, Rheumatoid Granulomatosis with Polyangiitis Leukemia Leukemia, Lymphoid Lymphoma Lymphoma, B-Cell Lymphoma, Follicular Lymphoma, Non-Hodgkin Myelitis Neuromyelitis Optica Purpura Purpura, Thrombocytopenic Purpura, Thrombocytopenic, Idiopathic Thrombocytopenia	-
Romidepsin	Lymphoma, T-Cell, Cutaneous	7
Rucaparib	Carcinoma, Ovarian Epithelial Fallopian Tube Neoplasms Peritoneal Neoplasms Prostatic Neoplasms, Castration-Resistant	54
Ruxolitinib	Graft vs Host Disease Polycythemia Polycythemia Vera Primary Myelofibrosis Thrombocytosis	6
Selinexor	Multiple Myeloma	83
Selumetinib	Neurofibromatosis 1	-
Siltuximab	Giant Lymph Node Hyperplasia	-
Sirolimus	Angiomyolipoma Constriction, Pathologic Coronary Restenosis Eye Diseases Immune System Diseases Kidney Failure, Chronic Lipoma Tuberous Sclerosis	86
Sonidegib	Carcinoma, Basal Cell	57
Sorafenib	Carcinoma, Hepatocellular Carcinoma, Renal Cell Thyroid Neoplasms	90
Sunitinib	Adenoma Carcinoma, Renal Cell Digestive System Neoplasms Gastrointestinal Neoplasms Gastrointestinal Stromal Tumors Intestinal Neoplasms	88
Talazoparib	Breast Neoplasms	59
Tamoxifen	Breast Diseases Cystic Fibrosis Cysts Fibroadenoma Fibrocystic Breast Disease Hemorrhage Menorrhagia Menstruation Disturbances Metrorrhagia Neoplasms	69
Tamsulosin	Calculi Coronary Artery Disease Heart Diseases Hernia Hernia, Inguinal Inflammation Ischemia Lithiasis Lower Urinary Tract Symptoms Myocardial Ischemia Prostatic Hyperplasia Ureteral Calculi Urinary Calculi Urolithiasis Urologic Diseases	-
Temozolomide	Astrocytoma Nervous System Neoplasms	55
Temsirolimus	Carcinoma, Renal Cell	83
Teniposide	Precursor Cell Lymphoblastic Leukemia-Lymphoma	77
Thalidomide	Brain Abscess Immune System Diseases Multiple Myeloma Neoplasms, Plasma Cell	-
Tivozanib	Carcinoma, Renal Cell	-
Tocilizumab	Arthritis Arthritis, Juvenile Arthritis, Rheumatoid Behavior Cytokine Release Syndrome Giant Cell Arteritis Neurobehavioral Manifestations Oral Manifestations Psychotic Disorders Schizophrenia Tic Disorders	-
Topotecan	Small Cell Lung Carcinoma	-
Toremifene	Breast Neoplasms	-
Trabectedin	Leiomyosarcoma Liposarcoma	-
Trametinib	Carcinoma, Non-Small-Cell Lung Melanoma	15
Trastuzumab	Breast Neoplasms Neoplasms	31
Tretinoin	Lentigo	92
Triptorelin	Fatty Liver Hypogonadism Infertility, Female Prostatic Neoplasms	8
Tucatinib	Breast Neoplasms	-
Valrubicin	Urinary Bladder Neoplasms	81
Vandetanib	Thyroid Neoplasms	94
Vemurafenib	Melanoma	4
Venetoclax	Leukemia, Lymphocytic, Chronic, B-Cell Leukemia, Myeloid, Acute	-
Vinblastine	Glioma	28
Vincristine	Precursor Cell Lymphoblastic Leukemia-Lymphoma	31
Vinorelbine	Carcinoma, Non-Small-Cell Lung	39
Vismodegib	Carcinoma, Basal Cell	57
Vorinostat	Lymphoma, T-Cell, Cutaneous	10
Zoledronate	Arthritis Bone Marrow Diseases Brain Abscess Chronic Kidney Disease-Mineral and Bone Disorder Chronic Periodontitis HIV	-

6. Conclusion

We applied the software package "Genome Enhancer" to a data set that contains *transcriptomics* data. The study is done in the context of *Squamous Cell Carcinoma*. The data were pre-processed, statistically analyzed and differentially expressed genes were identified. Also checked was the enrichment of GO or disease categories among the studied gene sets.

We propose the following drugs as most promising candidates for treating the pathology under study:



Erlotinib, Curcumin and {(2Z)-4-AMINO-2-[(4-METHOXYPHENYL)IMINO]-2,3-DIHYDRO-1,3-THIAZOL-5-YL}(4-METHOXYPHENYL)METHANONE

These drugs were selected for acting on the following targets: NTRK2, IGFBP5 and CCND1, which were predicted to be involved in the molecular mechanism of the pathology under study.

The identified molecular mechanism of the studied pathology was predicted to be mainly based on the following key drug targets:



integrins, ornithine decarboxylase, EGFR and EGFR

These potential drug targets should be considered as a prospective research initiative for further drug repurposing and drug development purposes. The following drugs were predicted as, matching those drug targets: 1-(5-OXO-2,3,5,9B-TETRAHYDRO-1H-PYRROLO[2,1-A]ISOINDOL-9-YL)-3-(5-PYRROLIDIN-2-YL-1H-PYRAZOL-3-YL)-UREA, Terlipressin, Cytarabine, Erlotinib, Tegafur, Curcumin, Methyl aminolevulinate and Eflornithine. These drugs should be considered with special caution for research purposes only.

In this study, we came up with a detailed signal transduction network regulating differentially expressed genes in the studied pathology. In this network we have revealed the following top master regulators (signaling proteins and their complexes) that play a crucial role in the molecular mechanism of the studied pathology, which can be proposed as the most promising molecular targets for further drug repurposing and drug development initiatives.

- integrins
- ornithine decarboxylase
- EGFR
- EGFR

Potential drug compounds which can be affecting these targets can be found in the "Finding prospective drug targets" section.

7. Methods

Databases used in the study

Transcription factor binding sites in promoters and enhancers of differentially expressed genes were analyzed using known DNA-binding motifs described in the **TRANSFAC**[®] library, release 2024.1 (geneXplain GmbH, Wolfenbüttel, Germany) (<https://genexplain.com/transfac>).

The master regulator search uses the **TRANSPATH**[®] database (BIOBASE), release 2024.1 (geneXplain GmbH, Wolfenbüttel, Germany) (<https://genexplain.com/transpath>). A comprehensive signal transduction network of human cells is built by the software on the basis of reactions annotated in **TRANSPATH**[®].

The information about drugs corresponding to identified drug targets and clinical trials references were extracted from **HumanPSD**[™] database, release 2024.1 (<https://genexplain.com/humanpsd>).

The Ensembl database release Human104.38 (hg38) (<http://www.ensembl.org>) was used for gene IDs representation and Gene Ontology (GO) (<http://geneontology.org>) was used for functional classification of the studied gene set.

Methods for the analysis of enriched transcription factor binding sites and composite modules

Transcription factor binding sites in promoters and enhancers of differentially expressed genes were analyzed using known DNA-binding motifs. The motifs are specified using position weight matrices (PWMs) that give weights to each nucleotide in each position of the DNA binding motif for a transcription factor or a group of them.

We search for transcription factor binding sites (TFBS) that are enriched in the promoters and enhancers under study as compared to a background sequence set such as promoters of genes that were not differentially regulated under the condition of the experiment. We denote study and background sets briefly as Yes and No sets. In the current work we used a workflow considering promoter sequences of a standard length of 1100 bp (-1000 to +100). The error rate in this part of the pipeline is controlled by estimating the adjusted p-value (using the Benjamini-Hochberg procedure) in comparison to the TFBS frequency found in randomly selected regions of the human genome (adj.p-value < 0.01).

We have applied the CMA algorithm (Composite Module Analyst) for searching composite modules [7] in the promoters and enhancers of the Yes and No sets. We searched for a composite module consisting of a cluster of 10 TFs in a sliding window of 200-300 bp that statistically significantly separates sequences in the Yes and No sets (minimizing Wilcoxon p-value).

Methods for finding master regulators in networks

We searched for master regulator molecules in signal transduction pathways upstream of the identified transcription factors. The master regulator search uses a comprehensive signal transduction network of human cells. The main algorithm of the master regulator search has been described earlier [3,4]. The goal of the algorithm is to find nodes in the global signal transduction network that may potentially regulate the activity of a set of transcription factors found at the previous step of the analysis. Such nodes are considered as most promising drug targets, since any influence on such a node may switch the transcriptional programs of hundreds of genes that are regulated by the respective TFs. In our analysis, we have run the algorithm with a maximum radius of 12 steps upstream of each TF in the input set. The error rate of this algorithm is controlled by applying it 10000 times to randomly generated sets of input transcription factors of the same set-size. Z-score and FDR value of ranks are calculated then for each potential master regulator node on the basis of such random runs (see detailed description in [9]). We control the error rate by the FDR threshold 0.05.

Methods for analysis of pharmaceutical compounds

We seek for the optimal combination of molecular targets (key elements of the regulatory network of the cell) that potentially interact with pharmaceutical compounds from a library of known drugs and biologically active chemical compounds, using information about known drugs from HumanPSD[™] and predicting potential drugs using **PASS** program.

Method for analysis of known pharmaceutical compounds

We selected compounds from HumanPSD™ database that have at least one target. Next, we sort compounds using "Drug rank" that is the sum of the following ranks:

1. ranking by "Target activity score" ($T\text{-score}_{PSD}$),
2. ranking by "Disease activity score" ($D\text{-score}_{PSD}$),
3. ranking by "Clinical validity score".

"Target activity score" ($T\text{-score}_{PSD}$) is calculated as follows:

$$T\text{-score}_{PSD} = -\frac{|T|}{|T| + w(|AT| - |T|)} \sum_{t \in T} \log_{10} \left(\frac{\text{rank}(t)}{1 + \max \text{Rank}(T)} \right),$$

where T is set of all targets related to the compound intersected with input list, $|T|$ is number of elements in T , AT and $|AT|$ are set set of all targets related to the compound and number of elements in it, w is weight multiplier, $\text{rank}(t)$ is rank of given target, $\max \text{Rank}(T)$ equals $\max(\text{rank}(t))$ for all targets t in T .

We use following formula to calculate "Disease activity score" ($D\text{-score}_{PSD}$):

$$D\text{-score}_{PSD} = \begin{cases} \sum_{d \in D} \sum_{p \in P} \text{phase}(d, p) \\ 0, D = \emptyset \end{cases},$$

where D is the set of selected diseases, and if D is empty set, $D\text{-score}_{PSD} = 0$. P is a set of all known phases for each disease, $\text{phase}(p, d)$ equals to the phase number if there are known clinical trials for the selected disease on this phase and zero otherwise.

The clinical validity score reflects the number of the highest clinical trials phase (from 1 to 4) on which the drug was ever tested for any pathology.

Method for prediction of pharmaceutical compounds

In this study, the focus was put on compounds with high pharmacological efficiency and low toxicity. For this purpose, comprehensive library of chemical compounds and drugs was subjected to a SAR/QSAR analysis. This library contains 13040 compounds along with their pre-calculated potential pharmacological activities of those substances, their possible side and toxic effects, as well as the possible mechanisms of action. All biological activities are expressed as probability values for a substance to exert this activity (Pa).

We selected compounds that satisfied the following conditions:

1. Toxicity below a chosen toxicity threshold (defines as Pa , probability to be active as toxic substance).
2. For all predicted pharmacological effects that correspond to a set of user selected disease(s) Pa is greater than a chosen effect threshold.
3. There are at least 2 targets (corresponding to the predicted activity-mechanisms) with predicted Pa greater than a chosen target threshold.

The maximum Pa value for all toxicities corresponding to the given compound is selected as the "Toxicity score". The maximum Pa value for all activities corresponding to the selected diseases for the given compound is used as the "Disease activity score". "Target activity score" (T-score) is calculated as follows:

$$T\text{-score}(s) = \frac{|T|}{|T| + w(|AT| - |T|)} \sum_{m \in M(s)} \left(pa(m) \sum_{g \in G(m)} IAP(g) \text{optWeight}(g) \right),$$

where $M(s)$ is the set of activity-mechanisms for the given structure (which passed the chosen threshold for activity-mechanisms Pa); $G(m)$ is the set of targets (converted to genes) that corresponds to the given activity-mechanism (m) for the given compound; $pa(m)$ is the probability to be active of the activity-mechanism (m), $IAP(g)$ is the invariant accuracy of prediction for gene from $G(m)$; $\text{optWeight}(g)$ is the additional weight multiplier for gene. T is set of all targets related to the compound intersected with input list, $|T|$ is number of elements in T , AT and $|AT|$ are set set of all targets related to the compound and number of elements in it, w is weight multiplier.

"Druggability score" (D-score) is calculated as follows:

$$D\text{-score}(g) = IAP(g) \sum_{s \in S(g)} \sum_{m \in M(s,g)} pa(m),$$

where $S(g)$ is the set of structures for which target list contains given target, $M(s,g)$ is the set of activity-mechanisms (for the given structure) that corresponds to the given gene, $pa(m)$ is the probability to be active of the activity-mechanism (m), $IAP(g)$ is the invariant accuracy of prediction for the given gene.

8. References

1. Kel A, Voss N, Jauregui R, Kel-Margoulis O, Wingender E. Beyond microarrays: Finding key transcription factors controlling signal transduction pathways. *BMC Bioinformatics*. **2006**;7(S2), S13. doi:10.1186/1471-2105-7-s2-s13
2. Stegmaier P, Voss N, Meier T, Kel A, Wingender E, Borlak J. Advanced Computational Biology Methods Identify Molecular Switches for Malignancy in an EGF Mouse Model of Liver Cancer. *PLoS ONE*. **2011**;6(3):e17738. doi:10.1371/journal.pone.0017738
3. Koschmann J, Bhar A, Stegmaier P, Kel A, Wingender E. “Upstream Analysis”: An Integrated Promoter-Pathway Analysis Approach to Causal Interpretation of Microarray Data. *Microarrays*. **2015**;4(2):270-286. doi:10.3390/microarrays4020270.
4. Kel A, Stegmaier P, Valeev T, Koschmann J, Poroikov V, Kel-Margoulis OV, and Wingender E. Multi-omics “upstream analysis” of regulatory genomic regions helps identifying targets against methotrexate resistance of colon cancer. *EuPA Open Proteom*. **2016**;13:1-13. doi:10.1016/j.euprot.2016.09.002
5. Michael H, Hogan J, Kel A et al. Building a knowledge base for systems pathology. *Brief Bioinformatics*. **2008**;9(6):518-531. doi:10.1093/bib/bbn038
6. Matys V, Kel-Margoulis OV, Fricke E, Liebich I, Land S, Barre-Dirrie A, Reuter I, Chekmenev D, Krull M, Hornischer K, Voss N, Stegmaier P, Lewicki-Potapov B, Saxel H, Kel AE, Wingender E. TRANSFAC and its module TRANSCompel: transcriptional gene regulation in eukaryotes. *Nucleic Acids Res*. **2006**;34(90001):D108-D110. doi:10.1093/nar/gkj143
7. Kel AE, Gössling E, Reuter I, Cheremushkin E, Kel-Margoulis OV, Wingender E. MATCH: A tool for searching transcription factor binding sites in DNA sequences. *Nucleic Acids Res*. **2003**;31(13):3576-3579. doi:10.1093/nar/gkg585
8. Waleev T, Shtokalo D, Konovalova T, Voss N, Cheremushkin E, Stegmaier P, Kel-Margoulis O, Wingender E, Kel A. Composite Module Analyst: identification of transcription factor binding site combinations using genetic algorithm. *Nucleic Acids Res*. **2006**;34(Web Server issue):W541-5.
9. Krull M, Pistor S, Voss N, Kel A, Reuter I, Kronenberg D, Michael H, Schwarzer K, Potapov A, Choi C, Kel-Margoulis O, Wingender E. TRANSPATH: an information resource for storing and visualizing signaling pathways and their pathological aberrations. *Nucleic Acids Res*. **2006**;34(90001):D546-D551. doi:10.1093/nar/gkj107
10. Boyarskikh U, Pintus S, Mandrik N, Stelmashenko D, Kiselev I, Evshin I, Sharipov R, Stegmaier P, Kolpakov F, Filipenko M, Kel A. Computational master-regulator search reveals mTOR and PI3K pathways responsible for low sensitivity of NCI-H292 and A427 lung cancer cell lines to cytotoxic action of p53 activator Nutlin-3. *BMC Med Genomics*. **2018**;11(1):12. doi:10.1186/1471-2105-7-s2-s13
11. Filimonov D, Poroikov V. Probabilistic Approaches in Activity Prediction. Varnek A, Tropsha A. *Cheminformatics Approaches to Virtual Screening*. Cambridge (UK): RSC Publishing. **2008**;:182-216.
12. Filimonov DA, Poroikov VV. Prognosis of specters of biological activity of organic molecules. *Russian chemical journal*. **2006**;50(2):66-75 (russ)
13. Filimonov D, Poroikov V, Borodina Y, Glorizova T. Chemical Similarity Assessment Through Multilevel Neighborhoods of Atoms: Definition and Comparison with the Other Descriptors. *ChemInform*. **1999**;39(4):666-670. doi:10.1002/chin.199940210

Thank you for using the Genome Enhancer!

In case of any questions please contact us at support@genexplain.com

Supplementary material

1. [Supplementary table 1 - Up-regulated genes](#)
2. [Supplementary table 2 - Down-regulated genes](#)
3. [Supplementary table 3 - Detailed report. Composite modules and master regulators \(up-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue\).](#)
4. [Supplementary table 4 - Detailed report. Composite modules and master regulators \(down-regulated genes in Experiment: Squamous Cell Carcinoma vs. Control: Non-tumour tissue\).](#)
5. [Supplementary table 5 - Detailed report. Pharmaceutical compounds and drug targets.](#)

Disclaimer

Decisions regarding care and treatment of patients should be fully made by attending doctors. The predicted chemical compounds listed in the report are given only for doctor's consideration and they cannot be treated as prescribed medication. It is the physician's responsibility to independently decide whether any, none or all of the predicted compounds can be used solely or in combination for patient treatment purposes, taking into account all applicable information regarding FDA prescribing recommendations for any therapeutic and the patient's condition, including, but not limited to, the patient's and family's medical history, physical examinations, information from various diagnostic tests, and patient preferences in accordance with the current standard of care. Whether or not a particular patient will benefit from a selected therapy is based on many factors and can vary significantly.

The compounds predicted to be active against the identified drug targets in the report are not guaranteed to be active against any particular patient's condition. GeneXplain GmbH does not give any assurances or guarantees regarding the treatment information and conclusions given in the report. There is no guarantee that any third party will provide a refund for any of the treatment decisions made based on these results. None of the listed compounds was checked by Genome Enhancer for adverse side-effects or even toxic effects.

The analysis report contains information about chemical drug compounds, clinical trials and disease biomarkers retrieved from the HumanPSD™ database of gene-disease assignments maintained and exclusively distributed worldwide by geneXplain GmbH. The information contained in this database is collected from scientific literature and public clinical trials resources. It is updated to the best of geneXplain's knowledge however we do not guarantee completeness and reliability of this information leaving the final checkup and consideration of the predicted therapies to the medical doctor.

The scientific analysis underlying the Genome Enhancer report employs a complex analysis pipeline which uses geneXplain's proprietary Upstream Analysis approach, integrated with TRANSFAC® and TRANSPATH® databases maintained and exclusively distributed worldwide by geneXplain GmbH. The pipeline and the databases are updated to the best of geneXplain's knowledge and belief, however, geneXplain GmbH shall not give a warranty as to the characteristics or to the content and any of the results produced by Genome Enhancer. Moreover, any warranty concerning the completeness, up-to-dateness, correctness and usability of Genome Enhancer information and results produced by it, shall be excluded.

The results produced by Genome Enhancer, including the analysis report, severely depend on the quality of input data used for the analysis. It is the responsibility of Genome Enhancer users to check the input data quality and parameters used for running the Genome Enhancer pipeline.

Note that the text given in the report is not unique and can be fully or partially repeated in other Genome Enhancer analysis reports, including reports of other users. This should be considered when publishing any results or excerpts from the report. This restriction refers only to the general description of analysis methods used for generating the report. All data and graphics referring to the concrete set of input data, including lists of mutated genes, differentially expressed genes/proteins/metabolites, functional classifications, identified transcription factors and master regulators, constructed molecular networks, lists of chemical compounds and reconstructed model of molecular mechanisms of the studied pathology are unique in respect to the used input data set and Genome Enhancer pipeline parameters used for the current run.